



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :  
A61K 39/00, 39/395, 48/00  
C07K 3/12, 15/06, 15/28  
C12N 15/12, C12P 19/34  
C12Q 1/68, G01N 33/53, 37/00

A1

(11) International Publication Number: WO 94/09820

(43) International Publication Date: 11 May 1994 (11.05.94)

(21) International Application Number: PCT/US93/10624

(22) International Filing Date: 5 November 1993 (05.11.93)

(30) Priority data:  
07/973,337 5 November 1992 (05.11.92) US

(60) Parent Application or Grant  
(63) Related by Continuation  
US --

07/973,337 (CIP)

(71) Applicant (for all designated States except US): SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH [US/US]; 1275 York Avenue, New York, NY 10021 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only) : ISRAELI, Ron, S. [US/US]; 950 49th Street, Apartment 6K, Brooklyn, NY 11219 (US). HESTON, Warren, D., W. [US/US]; 400 East 85th Street, Apartment 18B, New York, NY 10028 (US). FAIR, William, R. [US/US]; 400 East 70th Street, Apartment 3501, New York, NY 10021 (US).

(74) Agent: COBERT, Robert, J.; Cooper & Dunham, 30 Rockefeller Plaza, New York, NY 10112 (US).

(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

(54) Title: PROSTATE-SPECIFIC MEMBRANE ANTIGEN

(57) Abstract

This invention provides for an isolated mammalian nucleic acid molecule encoding a mammalian prostate-specific membrane antigen. This invention provides for nucleic acid probes which specifically hybridize with the nucleic acid molecule encoding said antigen. This invention provides for a method of detecting hematogenous micrometastatic tumor cells of a subject performing the polymerase chain reaction (PCR) on samples of the subject using primers of said antigen. This invention provides for methods to identify ligands which bind to said antigen. This invention provides for the prevention and/or treatment of prostate tumor growth.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PROSTATE-SPECIFIC MEMBRANE ANTIGEN

5

Background of the Invention

10 This application is a continuation-in-part of United States Application Serial No. 07/973,337, filed on November 5, 1992, the contents of which are hereby incorporated by reference.

15 This invention disclosed herein was made in part with Government support under NIH Grants No. DK47650 and CA58192 from the Department of Health and Human Services. Accordingly, the U.S. Government has certain rights in this invention.

20 Throughout this application various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.  
25 Full bibliographic citation for these references may be found at the end of of each series of experiments.

30 Prostate cancer is among the most significant medical problems in the United States, as the disease is now the most common malignancy diagnosed in American males. In 1992 there were over 132,000 new cases of prostate cancer detected with over 36,000 deaths attributable to the disease, representing a 17.3% increase over 4 years (2).  
35 Five year survival rates for patients with prostate cancer range from 88% for those with localized disease to 29% for those with metastatic disease. The rapid increase

-2-

in the number of cases appears to result in part from an increase in disease awareness as well as the widespread use of clinical markers such as the secreted proteins prostate-specific antigen (PSA) and prostatic acid phosphatase (PAP) (37).

The prostate gland is a site of significant pathology affected by conditions such as benign growth (BPH), neoplasia (prostatic cancer) and infection (prostatitis). Prostate cancer represents the second leading cause of death from cancer in man (1). However prostatic cancer is the leading site for cancer development in men. The difference between these two facts relates to prostatic cancer occurring with increasing frequency as men age, especially in the ages beyond 60 at a time when death from other factors often intervenes. Also, the spectrum of biologic aggressiveness of prostatic cancer is great, so that in some men following detection the tumor remains a latent histologic tumor and does not become clinically significant, whereas in other it progresses rapidly, metastasizes and kills the man in a relatively short 2-5 year period (1, 3).

In prostate cancer cells, two specific proteins that are made in very high concentrations are prostatic acid phosphatase (PAP) and prostate specific antigen (PSA) (4, 5, 6). These proteins have been characterized and have been used to follow response to therapy. With the development of cancer, the normal architecture of the gland becomes altered, including loss of the normal duct structure for the removal of secretions and thus the secretions reach the serum. Indeed measurement of serum PSA is suggested as a potential screening method for prostatic cancer. Indeed, the relative amount of PSA and/or PAP in the cancer reduces as compared to normal or

- 3 -

benign tissue.

PAP was one of the earliest serum markers for detecting metastatic spread (4). PAP hydrolyses tyrosine phosphate and has a broad substrate specificity. Tyrosine phosphorylation is often increased with oncogenic transformation. It has been hypothesized that during neoplastic transformation there is less phosphatase activity available to inactivate proteins that are activated by phosphorylation on tyrosine residues. In some instances, insertion of phosphatases that have tyrosine phosphatase activity has reversed the malignant phenotype.

PSA is a protease and it is not readily appreciated how loss of its activity correlates with cancer development (5, 6). The proteolytic activity of PSA is inhibited by zinc. Zinc concentrations are high in the normal prostate and reduced in prostatic cancer. Possibly the loss of zinc allows for increased proteolytic activity by PSA. As proteases are involved in metastasis and some proteases stimulate mitotic activity, the potentially increased activity of PSA could be hypothesized to play a role in the tumors metastases and spread (7).

Both PSA and PAP are found in prostatic secretions. Both appear to be dependent on the presence of androgens for their production and are substantially reduced following androgen deprivation.

Prostate-specific membrane antigen (PSM) which appears to be localized to the prostatic membrane has been identified. This antigen was identified as the result of generating monoclonal antibodies to a prostatic cancer cell, LNCaP (8).

- 4 -

Dr. Horoszewicz established a cell line designated LNCaP from the lymph node of a hormone refractory, heavily pretreated patient (9). This line was found to have an aneuploid human male karyotype. It maintained prostatic differentiation functionality in that it produced both PSA and PAP. It possessed an androgen receptor of high affinity and specificity. Mice were immunized with LNCaP cells and hybridomas were derived from sensitized animals. A monoclonal antibody was derived and was designated 7E11-C5 (8). The antibody staining was consistent with a membrane location and isolated fractions of LNCaP cell membranes exhibited a strongly positive reaction with immunoblotting and ELISA techniques. This antibody did not inhibit or enhance the growth of LNCaP cells in vitro or in vivo. The antibody to this antigen was remarkably specific to prostatic epithelial cells, as no reactivity was observed in any other component. Immunohistochemical staining of cancerous epithelial cells was more intense than that of normal or benign epithelial cells.

Dr. Horoszewicz also reported detection of immunoreactive material using 7E11-C5 in serum of prostatic cancer patients (8). The immunoreactivity was detectable in nearly 60% of patients with stage D-2 disease and in a slightly lower percentage of patients with earlier stage disease, but the numbers of patients in the latter group are small. Patients with benign prostatic hyperplasia (BPH) were negative. Patients with no apparent disease were negative, but 50-60% of patients in remission yet with active stable disease or with progression demonstrated positive serum reactivity. Patients with non prostatic tumors did not show immunoreactivity with 7E11-C5.

35

-5-

The 7E11-C5 monoclonal antibody is currently in clinical trials. The aldehyde groups of the antibody were oxidized and the linker-chelator glycol-tyrosyl- (n,  $\epsilon$ -diethylenetriamine-pentacetic acid)-lysine (GYK-DTPA) was coupled to the reactive aldehydes of the heavy chain (10). The resulting antibody was designated CYT-356. Immunohistochemical staining patterns were similar except that the CYT-356 modified antibody stained skeletal muscle. The comparison of CYT-356 with 7E11-C5 monoclonal antibody suggested both had binding to type 2 muscle fibers. The reason for the discrepancy with the earlier study, which reported skeletal muscle to be negative, was suggested to be due to differences in tissue fixation techniques. Still, the most intense and definite reaction was observed with prostatic epithelial cells, especially cancerous cells. Reactivity with mouse skeletal muscle was detected with immunohistochemistry but not in imaging studies. The Indium<sup>111</sup>-labeled antibody localized to LNCaP tumors grown in nude mice with an uptake of nearly 30% of the injected dose per gram tumor at four days. In-vivo, no selective retention of the antibody was observed in antigen negative tumors such as PC-3 and DU-145, or by skeletal muscle.

Very little was known about the PSM antigen. An effort at purification and characterization has been described at meetings by Dr. George Wright and colleagues (11, 12). These investigators have shown that following electrophoresis on acrylamide gels and Western blotting, the PSM antigen maintains a molecular weight of 100 kilodaltons (kd). Chemical and enzymatic treatment showed that both the peptide and carbohydrate moieties of the PSM antigen are required for recognition by the 7E11-C5 monoclonal antibody. Competitive binding studies with

-6-

specific lectins suggested that galNAc is the dominant carbohydrate of the antigenic epitope.

5 The 100kd glycoprotein unique to prostate cells and tissues was purified and characterized. The protein was digested proteolytically with trypsin and nine peptide fragments were sequenced. Using the technique of degenerate PCR (polymerase chain reaction), the full-length 2.65 kilobase (kb) cDNA coding for this antigen  
10 was cloned. Preliminary results have revealed that this antigen is highly expressed in prostate cancer tissues, including bone and lymph node metastases (13). The entire DNA sequence for the cDNA as well as the predicted amino acid sequence for the antigen was determined.  
15 Further characterization of the PSM antigen is presently underway in the applicants' laboratory including: analysis of PSM gene expression in a wide variety of tissues, transfection of the PSM gene into cells not expressing the antigen, chromosome localization of the  
20 PSM gene, cloning of the genomic PSM gene with analysis of the PSM promoter and generation of polyclonal and monoclonal antibodies against highly antigenic peptide domains of the PSM antigen, and identification of any endogenous PSM binding molecules (ligands).  
25  
Currently, LNCaP cells provide the best in-vitro model system to study human prostate cancer, since they produce all three prostatic bio-markers; PSA, PAP and PSM. The cells possess an aneuploid male karyotype with a Y  
30 chromosome, express a high affinity androgen receptor, and are hormonally responsive to both testosterone and DHT. Because PSM appears to be a transmembrane glycoprotein, it is considered an attractive target for both antibody-directed imaging and targeting of prostatic  
35 tumor deposits (38). We have demonstrated expression of



- 7 -

PSM protein in LNCaP cell membranes and in PC-3 cells transfected with PSM cDNA and also the characterization of PSM mRNA expression in human tissues, and in response to steroid hormones.

- 8 -

Brief Description of Figures

- Figure 1: Signal in lane 2 represent the 100kD PSM antigen. The EGFr was used as the positive control and is shown in lane 1. Incubation with rabbit antimouse (RAM) antibody alone served as negative control and is shown in lane 3.
- Figure 2 A-D: Upper two photos show LNCaP cytopins staining positively for PSM antigen. Lower left in DU-145 and lower right is PC-3 cytopsin, both negative for PSM antigen expression.
- Figure 3 A-D: Upper two panels are human prostate sections (BPH) staining positively for PSM antigen. The lower two panels show invasive prostate carcinoma human sections staining positively for expression of the PSM antigen.
- Figure 4: 100kD PSM antigen following immunoprecipitation of <sup>35</sup>S-Methionine labelled LNCaP cells with Cyt-356 antibody.
- Figure 5: 3% agarose gels stained with Ethidium bromide revealing PCR products obtained using the degenerate PSM antigen primers. The arrow points to sample IN-20, which is a 1.1 kb PCR product which we later confirmed to be a partial cDNA coding for

-9-

the PSM gene.

5                   **Figure 6 A-B:** 2% agarose gels of plasmid DNA resulting from TA cloning of PCR products. Inserts are excised from the PCR II vector (Invitrogen Corp.) by digestion with EcoRI. 1.1 kb PSM gene partial cDNA product is shown in lane 3 of gel 1.

10                   **Figure 7:** Autoradiogram showing size of cDNA represented in applicants' LNCaP library using M-MLV reverse transcriptase.

15                   **Figure 8:** Restriction analysis of full-length clones of PSM gene obtained after screening cDNA library. Samples have been cut with Not I and Sal I restriction enzymes to liberate the insert.

20                   **Figure 9:** Plasmid Southern autoradiogram of full length PSM gene clones. Size is approximately 2.7 kb.

25                   **Figure 10:** Northern blot revealing PSM expression limited to LNCaP prostate cancer line and H26 Ras-transfected LNCaP cell line. PC-3, DU-145, T-24, SKRC-27, HELA, MCF-7, HL-60, and others were are all negative.

30                   **Figure 11:** Autoradiogram of Northern analysis revealing expression of 2.8 kb PSM message unique to the LNCaP cell line (lane 1), and absent from the DU-145 (lane 2) and PC-3 cell lines (lane 3). RNA size ladder is shown on the left (kb), and 28S and 18S ribosomal RNA bands are indicated on the  
35

-10-

right.

5                   **Figure 12 A-B:** Results of PCR of human prostate tissues  
                    using PSM gene primers. Lanes are  
                    numbered from left to right. Lane 1,  
                    LNCaP; Lane 2, H26; Lane 3, DU-145; Lane  
                    4, Normal Prostate; Lane 5, BPH; Lane 6,  
                    Prostate Cancer; Lane 7, BPH; Lane 8,  
10                   Normal; Lane 9, BPH; Lane 10, BPH; Lane  
                    11, BPH; Lane 12, Normal; Lane 13, Normal;  
                    Lane 14, Cancer; Lane 15, Cancer; Lane 16,  
                    Cancer; Lane 17, Normal; Lane 18, Cancer;  
                    Lane 19, IN-20 Control; Lane 20, PSM cDNA

15           **Figure 13:**       Isoelectric point of PSM antigen (non-  
                                 glycosylated)

**Figure 14:1-8**   Secondary structure of PSM antigen

20           **Figure 15:A-B**   A. Hydrophilicity plot of PSM antigen  
                                 B. Prediction of membrane spanning  
                                 segments

**Figure 16:1-11** Homology with chicken, rat and human  
25                                   transferrin receptor sequence.

**Figure 17A-C:** Immunohistochemical detection of PSM  
                                 antigen expression in prostate cell lines.  
                                 Top panel reveals uniformly high level of  
30                                   expression in LNCaP cells; middle panel  
                                 and lower panel are DU-145 and PC-3 cells  
                                 respectively, both negative.

**Figure 18:**       Autoradiogram of protein gel revealing  
35                                   products of PSM coupled in-vitro

-11-

transcription/translation. Non-glycosylated PSM polypeptide is seen at 84 kDa (lane 1) and PSM glycoprotein synthesized following the addition of microsomes is seen at 100 kDa (lane 2).

**Figure 19:** Western Blot analysis detecting PSM expression in transfected non-PSM expressing PC-3 cells. 100 kDa PSM glycoprotein species is clearly seen in LNCaP membranes (lane 1), LNCaP crude lysate (lane 2), and PSM-transfected PC-3 cells (lane 4), but is undetectable in native PC-3 cells (lane 3).

**Figure 20:** Autoradiogram of ribonuclease protection gel assaying for PSM mRNA expression in normal human tissues. Radiolabeled 1 kb DNA ladder (Gibco-BRL) is shown in lane 1. Undigested probe is 400 nucleotides (lane 2), expected protected PSM band is 350 nucleotides, and tRNA control is shown (lane 3). A strong signal is seen in human prostate (lane 11), with very faint, but detectable signals seen in human brain (lane 4) and human salivary gland (lane 12).

**Figure 21:** Autoradiogram of ribonuclease protection gel assaying for PSM mRNA expression in LNCaP tumors grown in nude mice, and in human prostatic tissues. <sup>32</sup>P-labeled 1 kb DNA ladder is shown in lane 1. 298 nucleotide undigested probe is shown (lane

-12-

2), and tRNA control is shown (lane 3). PSM mRNA expression is clearly detectable in LNCaP cells (lane 4), orthotopically grown LNCaP tumors in nude mice with and without matrigel (lanes 5 and 6), and subcutaneously implanted and grown LNCaP tumors in nude mice (lane 7). PSM mRNA expression is also seen in normal human prostate (lane 8), and in a moderately differentiated human prostatic adenocarcinoma (lane 10). Very faint expression is seen in a sample of human prostate tissue with benign hyperplasia (lane 9).

15

**Figure 22:**

Ribonuclease protection assay for PSM expression in LNCaP cells treated with physiologic doses of various steroids for 24 hours. <sup>32</sup>P-labeled DNA ladder is shown in lane 1. 298 nucleotide undigested probe is shown (lane 2), and tRNA control is shown (lane 3). PSM mRNA expression is highest in untreated LNCaP cells in charcoal-stripped media (lane 4). Applicant see significantly diminished PSM expression in LNCaP cells treated with DHT (lane 5), Testosterone (lane 6), Estradiol (lane 7), and Progesterone (lane 8), with little response to Dexamethasone (lane 9).

30

**Figure 23:**

Data illustrating results of PSM DNA and RNA presence in transfect Dunning cell lines employing Southern and Northern

-13-

## blotting techniques

- 5      **Figure 24:A-B**      Figure A indicates the power of cytokine transfected cells to teach unmodified cells. Administration was directed to the parental flank or prostate cells. The results indicate the microenvironment considerations.
- 10           Figure B indicates actual potency at a particular site. The tumor was implanted in prostate cells and treated with immune cells at two different sites.
- 15      **Figure 25:A-B**      Relates potency of cytokines in inhibiting growth of primary tumors. Animals administered un-modified parental tumor cells and administered as a vaccine transfected cells. Following
- 20           prostatectomy of rodent tumor results in survival increase.
- 25      **Figure 26:**      PCR amplification with nested primers improved our level of detection of prostatic cells from approximately one prostatic cell per 10,000 MCF-7 cells to better than one cell per million MCF-7 cells, using either PSA.
- 30      **Figure 27:**      PCR amplification with nested primers improved our level of detection of prostatic cells from approximately one prostatic cell per 10,000 MCF-7 cells to better than one cell per million MCF-7
- 35           cells, using PSM-derived primers.

-14-

5      **Figure 28:**      A representative ethidium stained gel photograph for PSM-PCR. Samples run in lane A represent PCR products generated from the outer primers and samples in lanes labeled B are products of inner primer pairs.

10      **Figure 29:**      PSM Southern blot autoradiograph. The sensitivity of the Southern blot analysis exceeded that of ethidium staining, as can be seen in several samples where the outer product is not visible on figure 3 A-D, but is detectable by Southern blotting as shown in figure 4.

15      **Figure 30:**      Characteristics of the 16 patients analyzed with respect to their clinical stage, treatment, serum PSA and PAP values, and results of assay.



-15-

Summary of the Invention

This invention provides an isolated mammalian nucleic acid molecule encoding a mammalian prostate-specific membrane (PSM) antigen. The isolated mammalian nucleic acid may be DNA, cDNA or RNA.

This invention also provides nucleic acid molecule comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding the PSM antigen. The nucleic acid molecule may either be DNA or RNA.

This invention provides nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence of a nucleic acid molecule which is complementary to the nucleic acid molecule encoding a mammalian prostate-specific membrane antigen.

This invention further provides a method of detecting expression of the PSM antigen which comprises obtaining total mRNA from the cell and contacting the mRNA so obtained with a labelled PSM antigen specific nucleic acid molecule under hybridizing conditions, determining the presence of mRNA hybridized to the probe, and thereby detecting the expression of the PSM antigen by the cell. The PSM antigen in tissue sections may be similarly detected.

This invention provides isolated nucleic acid molecule of PSM antigen operatively linked to a promoter of RNA transcription. This invention further provides a vector which comprises an isolated mammalian nucleic acid molecule of PSM antigen.

-16-

This invention further provides a host vector system for the production of a polypeptide having the biological activity of a mammalian PSM antigen which comprises the vector comprising the mammalian nucleic acid molecule encoding a mammalian PSM antigen and a suitable host. The suitable host for the expression of PSM antigen may be a bacterial cell, insect cell, or mammalian cell.

This invention also provides a method of producing a polypeptide having the biological activity of a mammalian PSM antigen which comprises growing the host cell of vector system having a vector comprising the isolated mammalian nucleic acid molecule encoding a mammalian PSM antigen and a suitable host under suitable conditions permitting production of the polypeptide and recovery of the polypeptide so produced.

This invention provides a method for determining whether a ligand can bind to a mammalian PSM antigen which comprises contacting a mammalian cell having an isolated mammalian DNA molecule encoding a mammalian PSM antigen with the ligand under conditions permitting binding of ligands to the mammalian PSM antigen, and determining whether the ligand binds to a mammalian PSM antigen. This invention further provides ligands which bind to PSM antigen.

This invention provides purified mammalian PSM antigen. This invention also provides a polypeptide encoded by the isolated mammalian nucleic acid molecule encoding a mammalian PSM antigen. This invention further provides a method to identify and purify ligands of mammalian PSM antigen.

This invention further provides a method to produce both

-17-

polyclonal and monoclonal antibody using purified PSM antigens or polypeptides encoded by an isolated mammalian nucleic acid molecule encoding a mammalian PSM antigen.

5 This invention provides polyclonal and monoclonal antibody most likely but not limited to directed either to peptide Asp-Glu-Leu-Lys-Ala-Glu (SEQ ID No. 35), or Asn-Glu-Asp-Gly-Asn-Glu (SEQ ID No. 36) or Lys-Ser-Pro-Asp-Glu-Gly (SEQ ID No. 37) of the PSM antigen.

10 This invention provides a therapeutic agent comprising an antibody directed against a mammalian PSM antigen and a cytotoxic agent conjugated thereto.

15 This invention also provides a method of imaging prostate cancer in human patients which comprises administering to the patient at least one antibody directed against PSM antigen, capable of binding to the cell surface of the prostate cancer cell and labeled with an imaging agent  
20 under conditions so as to form a complex between the monoclonal antibody and the cell surface PSM antigen. This invention further provides a composition comprising an effective imaging amount of the antibody directed against PSM antigen and a pharmaceutically acceptable  
25 carrier.

This invention further provides a method of imaging prostate cancer in human patients which comprises administering to the patient multiple antibodies directed  
30 towards different PSM epitopes.

The invention also provides a method of imaging prostate cancer in human patients which comprises administering to the patient at least one ligand, capable of binding to  
35 the cell surface of the prostate cancer cell and labelled

-18-

with an imaging agent under conditions so as to form a complex between the ligand and the cell surface PSM antigen. This invention further provides a composition comprising an effective imaging amount of PSM antigen and  
5 a pharmaceutically acceptable carrier.

This invention provides an immunoassay for measuring the amount of the PSM antigen in a biological sample, e.g. serum, comprising steps of a) contacting the biological  
10 sample with at least one PSM antibody to form a complex with said antibody and the PSM antigen, and b) measuring the amount of PSM antigen in said biological sample by measuring the amount of said complex.

This invention also provides an immunoassay for measuring the amount of the PSM antigen in a biological sample comprising steps of a) contacting the biological sample with at least one PSM ligand to form a complex with said  
15 ligand and the PSM antigen, and b) measuring the amount of the PSM antigen in said biological sample by measuring the amount of said complex.  
20

This invention provides a method to purify mammalian PSM antigen comprising steps of: a) coupling the antibody  
25 directed against PSM antigen to a solid matrix; b) incubating the coupled antibody of a) with a cell lysate containing PSM antigen under the condition permitting binding of the antibody and PSM antigen; c) washing the coupled solid matrix to eliminate impurities and d)  
30 eluting the PSM antigen from the bound antibody.

This invention further provides transgenic nonhuman mammals which comprises an isolated nucleic acid molecule of PSM antigen. This invention also provides a  
35 transgenic nonhuman mammal whose genome comprises

-19-

antisense DNA complementary to DNA encoding a mammalian PSM antigen so placed as to be transcribed into antisense mRNA complementary to mRNA encoding the PSM antigen and which hybridizes to mRNA encoding the PSM antigen thereby  
5 reducing its translation.

This invention provides a method of suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor  
10 cells comprising introducing a DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor cell of a subject, in a way that expression of the prostate specific membrane antigen is under the control of the regulatory  
15 element, thereby suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells.

This invention provides a method of suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor  
20 cells, comprising introducing a DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element coupled with a therapeutic DNA  
25 into a tumor cell of a subject, thereby suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells.

This invention provides a therapeutic vaccine for preventing human prostate tumor growth or stimulation of prostate tumor cells in a subject, comprising administering an effective amount to the prostate cell, and a pharmaceutical acceptable carrier, thereby  
30 preventing the tumor growth or stimulation of tumor cells  
35

-20-

in the subject.

This invention provides a method of detecting hematogenous micrometastatic tumor cells of a subject, comprising (A) performing nested polymerase chain reaction (PCR) on blood, bone marrow, or lymph node samples of the subject using the prostate specific membrane antigen primers, and (B) verifying micrometastases by DNA sequencing and Southern analysis, thereby detecting hematogenous micrometastatic tumor cells of the subject.

This invention provides a method of abrogating the mitogenic response due to transferrin, comprising introducing a DNA molecule encoding prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor cell, the expression of which gene is directly associated with a defined pathological effect within a multicellular organism, thereby abrogating mitogen response due to transferrin.

-21-

Detailed Description of the Invention

Throughout this application, references to specific nucleotides are to nucleotides present on the coding strand of the nucleic acid. The following standard abbreviations are used throughout the specification to indicate specific nucleotides:

C=cytosine	A=adenosine
T=thymidine	G=guanosine

10

A "gene" means a nucleic acid molecule, the sequence of which includes all the information required for the normal regulated production of a particular protein, including the structural coding sequence, promoters and enhancers.

15

This invention provides an isolated mammalian nucleic acid encoding a mammalian prostate-specific membrane (PSM) antigen.

20

This invention further provides an isolated mammalian DNA molecule of an isolated mammalian nucleic acid molecule encoding a mammalian prostate-specific membrane antigen. This invention also provides an isolated mammalian cDNA molecule encoding a mammalian prostate-specific membrane antigen. This invention provides an isolated mammalian RNA molecule encoding a mammalian prostate-specific membrane antigen.

25

In the preferred embodiment of this invention, the isolated nucleic sequence is cDNA from human as shown in sequence ID number 1. This human sequence was submitted to GenBank (Los Alamos National Laboratory, Los Alamos, New Mexico) with Accession Number, M99487 and the description as PSM, Homo sapiens, 2653 base-pairs.

35

-22-

This invention also encompasses DNAs and cDNAs which encode amino acid sequences which differ from those of PSM antigen, but which should not produce phenotypic changes. Alternatively, this invention also encompasses  
5 DNAs and cDNAs which hybridize to the DNA and cDNA of the subject invention. Hybridization methods are well known to those of skill in the art.

The DNA molecules of the subject invention also include  
10 DNA molecules coding for polypeptide analogs, fragments or derivatives of antigenic polypeptides which differ from naturally-occurring forms in terms of the identity or location of one or more amino acid residues (deletion  
15 analogs containing less than all of the residues specified for the protein, substitution analogs wherein one or more residues specified are replaced by other residues and addition analogs where in one or more amino acid residues is added to a terminal or medial portion of the polypeptides) and which share some or all properties  
20 of naturally-occurring forms. These molecules include: the incorporation of codons "preferred" for expression by selected non-mammalian hosts; the provision of sites for cleavage by restriction endonuclease enzymes; and the provision of additional initial, terminal or intermediate  
25 DNA sequences that facilitate construction of readily expressed vectors.

The DNA molecules described and claimed herein are useful for the information which they provide concerning the  
30 amino acid sequence of the polypeptide and as products for the large scale synthesis of the polypeptide by a variety of recombinant techniques. The molecule is useful for generating new cloning and expression vectors, transformed and transfected prokaryotic and eukaryotic  
35 host cells, and new and useful methods for cultured



-23-

growth of such host cells capable of expression of the polypeptide and related products.

5 Moreover, the isolated mammalian nucleic acid molecules encoding a mammalian prostate-specific membrane antigen are useful for the development of probes to study the tumorigenesis of prostate cancer.

10 This invention also provides nucleic acid molecules of at least 15 nucleotides capable of specifically hybridizing with a sequence of a nucleic acid molecule encoding the prostate-specific membrane antigen.

15 This nucleic acid molecule produced can either be DNA or RNA. As used herein, the phrase "specifically hybridizing" means the ability of a nucleic acid molecule to recognize a nucleic acid sequence complementary to its own and to form double-helical segments through hydrogen bonding between complementary base pairs.

20 This nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence of a nucleic acid molecule encoding the prostate-specific membrane antigen can be used as a probe. Nucleic acid  
25 probe technology is well known to those skilled in the art who will readily appreciate that such probes may vary greatly in length and may be labeled with a detectable label, such as a radioisotope or fluorescent dye, to facilitate detection of the probe. DNA probe molecules  
30 may be produced by insertion of a DNA molecule which encodes PSM antigen into suitable vectors, such as plasmids or bacteriophages, followed by transforming into suitable bacterial host cells, replication in the transformed bacterial host cells and harvesting of the  
35 DNA probes, using methods well known in the art.

-24-

Alternatively, probes may be generated chemically from DNA synthesizers.

5 RNA probes may be generated by inserting the PSM antigen molecule downstream of a bacteriophage promoter such as T3, T7 or SP6. Large amounts of RNA probe may be produced by incubating the labeled nucleotides with the linearized PSM antigen fragment where it contains an upstream promoter in the presence of the appropriate RNA  
10 polymerase.

This invention also provides a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence of a nucleic acid molecule  
15 which is complementary to the mammalian nucleic acid molecule encoding a mammalian prostate-specific membrane antigen. This molecule may either be a DNA or RNA molecule.

20 The current invention further provides a method of detecting the expression of a mammalian PSM antigen expression in a cell which comprises obtaining total mRNA from the cell, contacting the mRNA so obtained with a labelled nucleic acid molecule of at least 15 nucleotides  
25 capable of specifically hybridizing with a sequence of the nucleic acid molecule encoding a mammalian PSM antigen under hybridizing conditions, determining the presence of mRNA hybridized to the molecule and thereby detecting the expression of the mammalian prostate-specific membrane antigen in the cell. The nucleic acid  
30 molecules synthesized above may be used to detect expression of a PSM antigen by detecting the presence of mRNA coding for the PSM antigen. Total mRNA from the cell may be isolated by many procedures well known to a  
35 person of ordinary skill in the art. The hybridizing

-25-

conditions of the labelled nucleic acid molecules may be determined by routine experimentation well known in the art. The presence of mRNA hybridized to the probe may be determined by gel electrophoresis or other methods known in the art. By measuring the amount of the hybrid made, the expression of the PSM antigen by the cell can be determined. The labelling may be radioactive. For an example, one or more radioactive nucleotides can be incorporated in the nucleic acid when it is made.

In one embodiment of this invention, nucleic acids are extracted by precipitation from lysed cells and the mRNA is isolated from the extract using an oligo-dT column which binds the poly-A tails of the mRNA molecules (13). The mRNA is then exposed to radioactively labelled probe on a nitrocellulose membrane, and the probe hybridizes to and thereby labels complementary mRNA sequences. Binding may be detected by luminescence autoradiography or scintillation counting. However, other methods for performing these steps are well known to those skilled in the art, and the discussion above is merely an example.

This invention further provides another method to detect expression of a PSM antigen in tissue sections which comprises contacting the tissue sections with a labelled nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence of nucleic acid molecules encoding a mammalian PSM antigen under hybridizing conditions, determining the presence of mRNA hybridized to the molecule and thereby detecting the expression of the mammalian PSM antigen in tissue sections. The probes are also useful for in-situ hybridization or in order to locate tissues which express this gene, or for other hybridization assays for the presence of this gene or its mRNA in various biological

-26-

tissues. The in-situ hybridization using a labelled nucleic acid molecule is well known in the art. Essentially, tissue sections are incubated with the labelled nucleic acid molecule to allow the hybridization to occur. The molecule will carry a marker for the detection because it is "labelled", the amount of the hybrid will be determined based on the detection of the amount of the marker and so will the expression of PSM antigen.

This invention further provides isolated PSM antigen nucleic acid molecule operatively linked to a promoter of RNA transcription. The isolated PSM antigen sequence can be linked to vector systems. Various vectors including plasmid vectors, cosmid vectors, bacteriophage vectors and other viruses are well known to ordinary skilled practitioners. This invention further provides a vector which comprises the isolated nucleic acid molecule encoding for the PSM antigen.

As an example to obtain these vectors, insert and vector DNA can both be exposed to a restriction enzyme to create complementary ends on both molecules which base pair with each other and are then ligated together with DNA ligase. Alternatively, linkers can be ligated to the insert DNA which correspond to a restriction site in the vector DNA, which is then digested with the restriction enzyme which cuts at that site. Other means are also available and known to an ordinary skilled practitioner.

In an embodiment, the PSM sequence is cloned in the Not I/Sal I site of pSPORT/vector (Gibco® - BRL). This plasmid, p55A-PSM, was deposited on August 14, 1992 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under

-27-

the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganism for the Purposes of Patent Procedure. Plasmid, p55A-PSM, was accorded ATCC Accession Number 75294.

5

This invention further provides a host vector system for the production of a polypeptide having the biological activity of the prostate-specific membrane antigen. These vectors may be transformed into a suitable host cell to form a host cell vector system for the production of a polypeptide having the biological activity of PSM antigen.

Regulatory elements required for expression include promoter sequences to bind RNA polymerase and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector includes a promoter such as the lac promoter and for transcription initiation the Shine-Dalgarno sequence and the start codon AUG (14). Similarly, a eukaryotic expression vector includes a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome. Such vectors may be obtained commercially or assembled from the sequences described by methods well known in the art, for example the methods described above for constructing vectors in general. Expression vectors are useful to produce cells that express the PSM antigen.

30

This invention further provides an isolated DNA or cDNA molecule described hereinabove wherein the host cell is selected from the group consisting of bacterial cells (such as E.coli), yeast cells, fungal cells, insect cells and animal cells. Suitable animal cells include, but are

35

-28-

not limited to Vero cells, HeLa cells, Cos cells, CV1 cells and various primary mammalian cells.

5 This invention further provides a method of producing a polypeptide having the biological activity of the prostate-specific membrane antigen which comprising growing host cells of a vector system containing the PSM antigen sequence under suitable conditions permitting production of the polypeptide and recovering the  
10 polypeptide so produced.

This invention provides a mammalian cell comprising a DNA molecule encoding a mammalian PSM antigen, such as a mammalian cell comprising a plasmid adapted for  
15 expression in a mammalian cell, which comprises a DNA molecule encoding a mammalian PSM antigen and the regulatory elements necessary for expression of the DNA in the mammalian cell so located relative to the DNA encoding the mammalian PSM antigen as to permit  
20 expression thereof.

Numerous mammalian cells may be used as hosts, including, but not limited to, the mouse fibroblast cell NIH3T3, CHO cells, HeLa cells, Ltk cells, Cos cells, etc. Expression  
25 plasmids such as that described supra may be used to transfect mammalian cells by methods well known in the art such as calcium phosphate precipitation, electroporation or DNA encoding the mammalian PSM antigen may be otherwise introduced into mammalian cells, e.g.,  
30 by microinjection, to obtain mammalian cells which comprise DNA, e.g., cDNA or a plasmid, encoding a mammalian PSM antigen.

This invention provides a method for determining whether

-29-

a ligand can bind to a mammalian prostate-specific membrane antigen which comprises contacting a mammalian cell comprising an isolated DNA molecule encoding a mammalian prostate-specific membrane antigen with the  
5 ligand under conditions permitting binding of ligands to the mammalian prostate-specific membrane antigen, and thereby determining whether the ligand binds to a mammalian prostate-specific membrane antigen.

10 This invention further provides ligands bound to the mammalian PSM antigen.

This invention also provides a therapeutic agent comprising a ligand identified by the above-described  
15 method and a cytotoxic agent conjugated thereto. The cytotoxic agent may either be a radioisotope or a toxin. Examples of radioisotopes or toxins are well known to one of ordinary skill in the art.

20 This invention also provides a method of imaging prostate cancer in human patients which comprises administering to the patients at least one ligand identified by the above-described method, capable of binding to the cell surface of the prostate cancer cell and labelled with an imaging  
25 agent under conditions permitting formation of a complex between the ligand and the cell surface PSM antigen. This invention further provides a composition comprising an effective imaging agent of the PSM antigen ligand and a pharmaceutically acceptable carrier. Pharmaceutically  
30 acceptable carriers are well known to one of ordinary skill in the art. For an example, such a pharmaceutically acceptable carrier can be physiological saline.

35 Also provided by this invention is a purified mammalian

-30-

PSM antigen. As used herein, the term "purified prostate-specific membrane antigen" shall mean isolated naturally-occurring prostate-specific membrane antigen or protein (purified from nature or manufactured such that the primary, secondary and tertiary conformation, and posttranslational modifications are identical to naturally-occurring material) as well as non-naturally occurring polypeptides having a primary structural conformation (i.e. continuous sequence of amino acid residues). Such polypeptides include derivatives and analogs.

This invention further provides a polypeptide encoded by the isolated mammalian nucleic acid sequence of PSM antigen.

It is believed that there may be natural ligand interacting with the PSM antigen. This invention provides a method to identify such natural ligand or other ligand which can bind to the PSM antigen. A method to identify the ligand comprises a) coupling the purified mammalian PSM antigen to a solid matrix, b) incubating the coupled purified mammalian PSM protein with the potential ligands under the conditions permitting binding of ligands and the purified PSM antigen; c) washing the ligand and coupled purified mammalian PSM antigen complex formed in b) to eliminate the nonspecific binding and impurities and finally d) eluting the ligand from the bound purified mammalian PSM antigen. The techniques of coupling proteins to a solid matrix are well known in the art. Potential ligands may either be deduced from the structure of mammalian PSM or by other empirical experiments known by ordinary skilled practitioners. The conditions for binding may also easily be determined and protocols for carrying such experimentation have long



- 31 -

been well documented (15). The ligand-PSM antigen complex will be washed. Finally, the bound ligand will be eluted and characterized. Standard ligands characterization techniques are well known in the art.

5

The above method may also be used to purify ligands from any biological source. For purification of natural ligands in the cell, cell lysates, serum or other biological samples will be used to incubate with the  
10 mammalian PSM antigen bound on a matrix. Specific natural ligand will then be identified and purified as above described.

15

With the protein sequence information, antigenic areas may be identified and antibodies directed against these areas may be generated and targeted to the prostate cancer for imaging the cancer or therapies.

20

This invention provides an antibody directed against the amino acid sequence of a mammalian PSM antigen.

25

This invention provides a method to select specific regions on the PSM antigen to generate antibodies. The protein sequence may be determined from the PSM DNA sequence. Amino acid sequences may be analyzed by methods well known to those skilled in the art to determine whether they produce hydrophobic or hydrophilic regions in the proteins which they build. In the case of cell membrane proteins, hydrophobic regions are well  
30 known to form the part of the protein that is inserted into the lipid bilayer of the cell membrane, while hydrophilic regions are located on the cell surface, in an aqueous environment. Usually, the hydrophilic regions will be more immunogenic than the hydrophobic regions.  
35 Therefore the hydrophilic amino acid sequences may be

- 32 -

selected and used to generate antibodies specific to mammalian PSM antigen. For an example, hydrophilic sequences of the human PSM antigen shown in hydrophilicity plot of Figure 16 may be easily selected.

5 The selected peptides may be prepared using commercially available machines. As an alternative, DNA, such as a cDNA or a fragment thereof, may be cloned and expressed and the resulting polypeptide recovered and used as an immunogen.

10 Polyclonal antibodies against these peptides may be produced by immunizing animals using the selected peptides. Monoclonal antibodies are prepared using hybridoma technology by fusing antibody producing B cells  
15 from immunized animals with myeloma cells and selecting the resulting hybridoma cell line producing the desired antibody. Alternatively, monoclonal antibodies may be produced by in vitro techniques known to a person of ordinary skill in the art. These antibodies are useful  
20 to detect the expression of mammalian PSM antigen in living animals, in humans, or in biological tissues or fluids isolated from animals or humans.

In one embodiment, peptides Asp-Glu-Leu-Lys-Ala-Glu (SEQ  
25 ID No. 35), Asn-Glu-Asp-Gly-Asn-Glu (SEQ ID No. 36) and Lys-Ser-Pro-Asp-Glu-Gly (SEQ ID No. 37) of human PSM antigen are selected.

This invention further provides polyclonal and monoclonal  
30 antibody(ies) against peptides Asp-Glu-Leu-Lys-Ala-Glu (SEQ ID No. 35), Asn-Glu-Asp-Gly-Asn-Glu (SEQ ID No. 36) and Lys-Ser-Pro-Asp-Glu-Gly (SEQ ID No. 37).

This invention provides a therapeutic agent comprising  
35 antibodies or ligand(s) directed against PSM antigen and

- 33 -

a cytotoxic agent conjugated thereto or antibodies linked enzymes which activate prodrug to kill the tumor. The cytotoxic agent may either be a radioisotope or toxin.

- 5 This invention provides a method of imaging prostate cancer in human patients which comprises administering to the patient the monoclonal antibody directed against the peptide of the mammalian PSM antigen capable of binding to the cell surface of the prostate cancer cell and
- 10 labeled with an imaging agent under conditions permitting formation of a complex between the monoclonal antibody and the cell surface prostate-specific membrane antigen. The imaging agent is a radioisotope such as Indium<sup>111</sup>.
- 15 This invention further provides a prostate cancer specific imaging agent comprising the antibody directed against PSM antigen and a radioisotope conjugated thereto.
- 20 This invention also provides a composition comprising an effective imaging amount of the antibody directed against the PSM antigen and a pharmaceutically acceptable carrier. The methods to determine effective imaging amounts are well known to a skilled practitioner. One
- 25 method is by titration using different amounts of the antibody.

- This invention further provides an immunoassay for measuring the amount of the prostate-specific membrane antigen in a biological sample comprising steps of a)
- 30 contacting the biological sample with at least one antibody directed against the PSM antigen to form a complex with said antibody and the prostate-specific membrane antigen, and b) measuring the amount of the

- 34 -

prostate-specific membrane antigen in said biological sample by measuring the amount of said complex. One example of the biological sample is a serum sample.

5 This invention provides a method to purify mammalian prostate-specific membrane antigen comprising steps of a) coupling the antibody directed against the PSM antigen to a solid matrix; b) incubating the coupled antibody of a) with lysate containing prostate-specific membrane antigen  
10 under the condition which the antibody and prostate membrane specific can bind; c) washing the solid matrix to eliminate impurities and d) eluting the prostate-specific membrane antigen from the coupled antibody.

15 This invention also provides a transgenic nonhuman mammal which comprises the isolated nucleic acid molecule encoding a mammalian PSM antigen. This invention further provides a transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a  
20 mammalian prostate-specific membrane antigen so placed as to be transcribed into antisense mRNA complementary to mRNA encoding the prostate-specific membrane antigen and which hybridizes to mRNA encoding the prostate specific antigen thereby reducing its translation.

25 Animal model systems which elucidate the physiological and behavioral roles of mammalian PSM antigen are produced by creating transgenic animals in which the expression of the PSM antigen is either increased or  
30 decreased, or the amino acid sequence of the expressed PSM antigen is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a mammalian PSM antigen, by microinjection,  
35 electroporation, retroviral transfection or other means

-35-

well known to those skilled in the art, into appropriate fertilized embryos in order to produce a transgenic animal (16) or 2) Homologous recombination (17) of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these PSM antigen sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and so is useful for producing an animal that cannot express native PSM antigen but does express, for example, an inserted mutant PSM antigen, which has replaced the native PSM antigen in the animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and so is useful for producing an animal which expresses its own and added PSM antigens, resulting in overexpression of the PSM antigens.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as M2 medium (16). DNA or cDNA encoding a mammalian PSM antigen is purified from a vector by methods well known in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the trans-gene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the trans-gene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a pipet puller) and the egg to be injected is put in a depression slide. The needle is

-36-

inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse (a mouse stimulated by the appropriate hormones to maintain pregnancy but which is not actually pregnant), where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg cell, and is used here only for exemplary purposes.

Another use of the PSM antigen sequence is to isolate homologous gene or genes in different mammals. The gene or genes can be isolated by low stringency screening of either cDNA or genomic libraries of different mammals using probes from PSM sequence. The positive clones identified will be further analyzed by DNA sequencing techniques which are well known to an ordinary person skilled in the art. For example, the detection of members of the protein serine kinase family by homology probing (18).

This invention provides a method of suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells comprising introducing a DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor cell of a subject, in a way that expression of the prostate specific membrane antigen is under the control of the regulatory element, thereby suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells. The subject may be a mammal or more specifically a human.

In one embodiment, the DNA molecule encoding prostate

-37-

specific membrane antigen operatively linked to a 5' regulatory element forms part of a transfer vector which is inserted into a cell or organism. In addition the vector is capable of replication and expression of prostate specific membrane antigen. The DNA molecule encoding prostate specific membrane antigen can be integrated into a genome of a eukaryotic or prokaryotic cell or in a host cell containing and/or expressing a prostate specific membrane antigen.

Further, the DNA molecule encoding prostate specific membrane antigen may be introduced by a bacterial, viral, fungal, animal, or liposomal delivery vehicle. Other means are also available and known to an ordinary skilled practitioner.

Further, the DNA molecule encoding a prostate specific membrane antigen operatively linked to a promoter or enhancer. A number of viral vectors have been described including those made from various promoters and other regulatory elements derived from virus sources. Promoters consist of short arrays of nucleic acid sequences that interact specifically with cellular proteins involved in transcription. The combination of different recognition sequences and the cellular concentration of the cognate transcription factors determines the efficiency with which a gene is transcribed in a particular cell type.

Examples of suitable promoters include a viral promoter. Viral promoters include: adenovirus promoter, an simian virus 40 (SV40) promoter, a cytomegalovirus (CMV) promoter, a mouse mammary tumor virus (MMTV) promoter, a Malony murine leukemia virus promoter, a murine sarcoma virus promoter, and a Rous sarcoma virus promoter.

-38-

Further, another suitable promoter is a heat shock promoter. Additionally, a suitable promoter is a bacteriophage promoter. Examples of suitable bacteriophage promoters include but not limited to, a T7 promoter, a T3 promoter, an SP6 promoter, a lambda promoter, a baculovirus promoter.

Also suitable as a promoter is an animal cell promoter such as an interferon promoter, a metallothionein promoter, an immunoglobulin promoter. A fungal promoter is also a suitable promoter. Examples of fungal promoters include but are not limited to, an ADC1 promoter, an ARG promoter, an ADH promoter, a CYC1 promoter, a CUP promoter, an ENO1 promoter, a GAL promoter, a PHO promoter, a PGK promoter, a GAPDH promoter, a mating type factor promoter. Further, plant cell promoters and insect cell promoters are also suitable for the methods described herein.

This invention provides a method of suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells, comprising introducing a DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element coupled with a therapeutic DNA into a tumor cell of a subject, thereby suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells. The subject may be a mammal or more specifically a human.

Further, the therapeutic DNA which is coupled to the DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor cell may code for a cytokine, viral antigen, or a pro-drug activating enzyme. Other means are also available and known



-39-

to an ordinary skilled practitioner.

5 The cytokine used may be interleukin-2, interleukin-12, interferon alpha, beta or gamma, granulocytic macrophage - colony stimulating factor, or other immunity factors.

10 In addition, this invention provides a prostate tumor cell, comprising a DNA molecule isolated from mammalian nucleic acid encoding a mammalian prostate-specific membrane antigen under the control of a prostate specific membrane antigen operatively linked to a 5' regulatory element.

15 As used herein, DNA molecules include complementary DNA (cDNA), synthetic DNA, and genomic DNA.

20 This invention provides a therapeutic vaccine for preventing human prostate tumor growth or stimulation of prostate tumor cells in a subject, comprising administering an effective amount to the prostate cell, and a pharmaceutical acceptable carrier, thereby preventing the tumor growth or stimulation of tumor cells in the subject. Other means are also available and known to an ordinary skilled practitioner.

25

This invention provides a method of detecting hematogenous micrometastatic tumor cells of a subject, comprising (A) performing nested polymerase chain reaction (PCR) on blood, bone marrow or lymph node samples of the subject using the prostate specific membrane antigen primers, and (B) verifying micrometastases by DNA sequencing and Southern analysis, thereby detecting hematogenous micrometastatic tumor cells of the subject. The subject may be a mammal or more specifically a human.

35

The micrometastatic tumor cell may be a prostatic cancer

-40-

and the DNA primers may be derived from prostate specific antigen. Further, the subject may be administered with simultaneously an effective amount of hormones, so as to increase expression of prostate specific membrane antigen.

5

This invention provides a method of abrogating the mitogenic response due to transferrin, comprising introducing a DNA molecule encoding prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor  
10 cell, the expression of which gene is directly associated with a defined pathological effect within a multicellular organism, thereby abrogating mitogen response due to transferrin. The tumor cell may be a prostate cell.

15 This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow  
20 thereafter.

## Experimental Details

### First Series of Experiments

#### 5 Materials and Methods

The approach for cloning the gene involved purification of the antigen in large quantities by immunoprecipitation, and microsequencing of several internal peptides for use in synthesizing degenerate oligonucleotide primers for subsequent use in the polymerase chain reaction (19, 20). A partial cDNA was amplified as a PCR product and this was used as a homologous probe to clone the full-length cDNA molecule from a LNCaP (Lymph Node Carcinoma of Prostate) cell line cDNA plasmid library (8). Early experiments revealed to us that the CYT-356 antibody (9) was not capable of detecting the antigen produced in bacteria since the epitope was the glycosylated portion of the PSM antigen, and this necessitated our more difficult, yet elaborate approach.

20

#### Western Analysis of the PSM Antigen

Membrane proteins were isolated from cells by hypotonic lysis followed by centrifugation over a sucrose density gradient (21). 10-20 $\mu$ g of LNCaP, DU-145, and PC-3 membrane proteins were electrophoresed through a 10% SDS-PAGE resolving gel with a 4% stacking gel at 9-10 milliamps for 16-18 hours. Proteins were electroblotted onto PVDF membranes (Millipore<sup>®</sup> Corp.) in transfer buffer (48mM Tris base, 39mM Glycine, 20% Methanol) at 25 volts overnight at 4°C. Membranes were blocked in TSB (0.15M NaCl, 0.01M Tris base, 5% BSA) for 30 minutes at room temperature followed by incubation with 10-15 $\mu$ g/ml of CYT-356 monoclonal antibody (Cytogen Corp.) for 2 hours. Membranes were then incubated with 10-15 $\mu$ g/ml of rabbit anti-mouse immunoglobulin (Accurate Scientific) for 1 hour at room temperature

-42-

followed by incubation with  $^{125}\text{I}$ -Protein A (Amersham®) at  $1 \times 10^6$  cpm/ml at room temperature. Membranes were then washed and autoradiographed for 12-24 hours at  $-70^\circ\text{C}$  (Figure 1).

5

#### Immunohistochemical Analysis of PSM Antigen Expression

The avidin-biotin method of immunohistochemical detection was employed to analyze both human tissue sections and cell lines for PSM Antigen expression (22). Cryostat-cut prostate tissue sections (4-6  $\mu\text{m}$  thick) were fixed in methanol/acetone for 10 minutes. Cell cytopspins were made on glass slides using 50,000 cells/100  $\mu\text{l}$ /slide. Samples were treated with 1% hydrogen peroxide in PBS for 10-15 minutes in order to remove any endogenous peroxidase activity. Tissue sections were washed several times in PBS, and then incubated with the appropriate suppressor serum for 20 minutes. The suppressor serum was drained off and the sections or cells were then incubated with the diluted CYT-356 monoclonal antibody for 1 hour. Samples were then washed with PBS and sequentially incubated with secondary antibodies (horse or goat immunoglobulins, 1:200 dilution for 30 minutes), and with avidin-biotin complexes (1:25 dilution for 30 minutes). DAB was used as a chromogen, followed by hematoxylin counterstaining and mounting. Frozen sections of prostate samples and duplicate cell cytopspins were used as controls for each experiment. As a positive control, the anti-cytokeratin monoclonal antibody CAM 5.2 was used following the same procedure described above. Tissue sections are considered by us to express the PSM antigen if at least 5% of the cells demonstrate immunoreactivity. Our scoring system is as follows: 1 = <5%; 2 = 5-19%; 3 = 20-75%; and 4 = >75% positive cells. Homogeneity versus heterogeneity was accounted for by evaluating positive and negative cells in 3-5 high power

-43-

light microscopic fields (400x), recording the percentage of positive cells among 100-500 cells. The intensity of immunostaining is graded on a 1+ to 4+ scale, where 1+ represents mild, 2-3+ represents moderate, and 4+ represents  
5 intense immunostaining as compared to positive controls.

#### Immunoprecipitation of the PSM Antigen

80%-confluent LNCaP cells in 100mm petri dishes were starved in RPMI media without methionine for 2 hours, after which  
10 <sup>35</sup>S-Methionine was added at 100μCi/ml and the cells were grown for another 16-18 hours. Cells were then washed and lysed by the addition of 1ml of lysis buffer (1% Triton X-100, 50mM Hepes pH 7.5, 10% glycerol, 150mM MgCl<sub>2</sub>, 1mM PMSF, and 1mM EGTA) with incubation for 20 minutes at 4°C.  
15 Lysates were pre-cleared by mixing with Pansorbin® cells (Calbiochem®) for 90 minutes at 4°C. Cell lysates were then mixed with Protein A Sepharose® CL-4B beads (Pharmacia®) previously bound with CYT-356 antibody (Cytogen Corp.) and RAM antibody (Accurate Scientific) for 3-4 hours at 4°C.  
20 12μg of antibody was used per 3mg of beads per petri dish. Beads were then washed with HNTG buffer (20mM Hepes pH 7.5, 150mM NaCl, 0.1% Triton X-100, 10% glycerol, and 2mM Sodium Orthovanadate), resuspended in sample loading buffer containing β-mercaptoethanol, denatured at 95°C for 5-10  
25 minutes and run on a 10% SDS-PAGE gel with a 4° stacking gel at 10 milliamps overnight. Gels were stained with Coomassie Blue, destained with acetic acid/methanol, and dried down in a vacuum dryer at 60°C. Gels were then autoradiographed for 16-24 hours at -70°C (Figure 2 A-D).

30

#### Large-Scale Immunoprecipitation and Peptide Sequencing

The procedure described above for immunoprecipitation was repeated with 8 confluent petri dishes containing approximately 6x10<sup>7</sup> LNCaP cells. The immunoprecipitation

- 44 -

product was pooled and loaded into two lanes of a 10% SDS-PAGE gel and electrophoresed at 9-10 milliamps for 16 hours. Proteins were electroblotted onto Nitrocellulose BA-85 membranes (Schleicher and Schuell®) for 2 hours at 75 volts at 4°C in transfer buffer. Membranes were stained with Ponceau Red to visualize the proteins and the 100kD protein band was excised, solubilized, and digested proteolytically with trypsin. HPLC was then performed on the digested sample on an Applied Biosystems Model 171C and clear dominant peptide peaks were selected and sequenced by modified Edman degradation on a modified post liquid Applied Biosystems Model 477A Protein/Peptide Microsequencer (23). Sequencing data on all of the peptides is included within this document. We attempted to sequence the amino-terminus of the PSM antigen by a similar method which involved purifying the antigen by immunoprecipitation and transfer via electro-blotting to a PVDF membrane (Millipore®). Protein was analyzed on an Applied Biosystems Model 477A Protein/Peptide Sequencer and the amino terminus was found to be blocked, and therefore no sequence data could be obtained by this technique.

## PSM Antigen Peptide Sequences:

25	2T17 #5	SLYES(W)TK (SEQ ID No. 3)
	2T22 #9	(S)YPDGXNLP GG(g)VQR (SEQ ID No. 4)
	2T26 #3	FYDPMFK (SEQ ID No. 5)
	2T27 #4	IYNVIGTL(K) (SEQ ID No. 6)
	2T34 #6	FLYXXTQIPHLAGTEQNFQLAK (SEQ ID No. 7)
30	2T35 #2	G/PVILYSDPADYFAPD/GVK (SEQ ID No. 8, 9)
	2T38 #1	AFIDPLGLPDRPFYR (SEQ ID No. 10)
	2T46 #8	YAGESFPGIYDALFDIESK (SEQ ID No. 11)
	2T47 #7	TILFAS(W)DAEEFGXX(q)STE(e)A(E)..(SEQ ID No. 12)

35 Notes: X means that no residue could be identified at this

-45-

position. Capital denotes identification but with a lower degree of confidence. (lower case) means residue present but at very low levels. ... indicates sequence continues but has dropped below detection limit.

5

All of these peptide sequences were verified to be unique after a complete homology search of the translated Genbank computer database.

10 Degenerate PCR

Sense and anti-sense 5'-unphosphorylated degenerate oligonucleotide primers 17 to 20 nucleotides in length corresponding to portions of the above peptides were synthesized on an Applied Biosystems Model 394A DNA  
15 Synthesizer. These primers have degeneracies from 32 to 144. The primers used are shown below. The underlined amino acids in the peptides represent the residues used in primer design.

20 Peptide 3: FYDPMFK (SEQ ID No. 5)

PSM Primer "A" TT(C or T) - TA(C or T) - GA(C or T) - CCX -  
ATG - TT (SEQ ID No. 13)

25 PSM Primer "B" AAC - ATX - GG(A or G) - TC(A or G) - TA(A  
or G) - AA (SEQ ID No. 14)

Primer A is sense primer and B is anti-sense. Degeneracy is 32-fold.

30

Peptide 4: IYNVIGTL(K) (SEQ ID No. 6)

PSM Primer "C" AT(T or C or A) - TA(T or C) - AA(T or C) -  
35 GTX - AT(T or C or A) - GG (SEQ ID No. 15)

-46-

PSM Primer "D" CC(A or T or G) - ATX - AC(G or A) - TT(A or G) - TA(A or G or T) - AT (SEQ ID No. 16)

Primer C is sense primer and D is anti-sense. Degeneracy is  
5 144-fold.

Peptide 2: G/PVILYSDPADYFAPD/GVK (SEQ ID No. 8,9)

PSM Primer "E" CCX - GCX - GA(T or C) - TA(T or C) - TT(T  
10 or C) - GC (SEQ ID No. 17)

PSM Primer "F" GC(G or A) - AA(A or G) - TA(A or G) - TXC -  
GCX - GG (SEQ ID No. 18)

15 Primer E is sense primer and F is antisense primer.  
Degeneracy is 128-fold.

Peptide 6: FLYXXTQIPHLAGTEQNFQLAK (SEQ ID No. 7)

20 PSM Primer "I" ACX - GA(A or G) - CA(A or G) - AA(T or C) -  
TT(T or C) - CA(A or G) - CT (SEQ ID No. 19)

PSM Primer "J" AG - (T or C)TG - (A or G)AA - (A or G)TT -  
(T or C)TG - (T or C)TC - XGT (SEQ ID No. 20)

25

PSM Primer "K" GA(A or G) - CA(A or G) - AA(T or C) - TT(T  
or C) CA(A or G) - CT (SEQ ID No. 21)

PSM Primer "L" AG - (T or C)TG - (A or G)AA - (A or G)TT -  
30 (T or C)TG - (T or C)TC (SEQ ID No. 22)

Primers I and K are sense primers and J and L are anti-  
sense. I and J have degeneracies of 128-fold and K and L  
have 32-fold degeneracy.

35



-47-

Peptide 7:        TILFAS(W)DAEEFGXX(q)STE(e)A(E)... (SEQ ID No. 12)

PSM Primer "M"    TGG - GA(T or C) - GCX - GA(A or G) - GA(A  
5   or G) - TT(C or T) - GG (SEQ ID No. 23)

PSM Primer "N"    CC - (G or A)AA - (T or C)TC - (T or C)TC -  
XGC - (A or G)TC - CCA (SEQ ID No. 24)

10   PSM Primer "O"   TGG - GA(T or C) - GCX - GA(A or G) - GA(A  
or G) - TT (SEQ ID No. 25)

PSM Primer "P"    AA - (T or C)TC - (T or C)TC - XGC - (A or  
G)TC - CCA (SEQ ID No. 26)

15

Primers M and O are sense primers and N and P are anti-sense. M and N have degeneracy of 64-fold and O and P are 32-fold degenerate.

20   Degenerate PCR was performed using a Perkin-Elmer Model 480  
DNA thermal cycler. cDNA template for the PCR was prepared  
from LNCaP mRNA which had been isolated by standard methods  
of oligo dT chromatography (Collaborative Research). The  
cDNA synthesis was carried out as follows:

25

4.5 $\mu$ l        LNCaP poly A+ RNA (2 $\mu$ g)  
1.0 $\mu$ l        Oligo dT primers (0.5 $\mu$ g)  
4.5 $\mu$ l        dH<sub>2</sub>O  
10 $\mu$ l

30

Incubate at 68°C x 10 minutes.  
Quick chill on ice x 5 minutes.

Add:

- 48 -

4 $\mu$ l        5 x RT Buffer  
2 $\mu$ l        0.1M DTT  
1 $\mu$ l        10mM dNTPs  
0.5 $\mu$ l      RNasin (Promega)  
5    1.5 $\mu$ l      dH<sub>2</sub>O  
19 $\mu$ l

Incubate for 2 minutes at 37°C.

Add 1 $\mu$ l Superscript® Reverse Transcriptase (Gibco®-BRL)

10    Incubate for 1 hour at 37°C.

Add 30 $\mu$ l dH<sub>2</sub>O.

Use 2 $\mu$ l per PCR reaction.

15    Degenerate PCR reactions were optimized by varying the  
annealing temperatures, Mg++ concentrations, primer  
concentrations, buffer composition, extension times and  
number of cycles. Our optimal thermal cycler profile was:  
Denaturation at 94°C x 30 seconds, Annealing at 45-55°C for  
20    1 minute (depending on the mean T<sub>m</sub> of the primers used), and  
Extension at 72°C for 2 minutes.

5 $\mu$ l        10 x PCR Buffer\*  
5 $\mu$ l        2.5mM dNTP Mix  
25    5 $\mu$ l        Primer Mix (containing 0.5-1.0 $\mu$ g each of sense &  
anti-sense primers)  
5 $\mu$ l        100mM  $\beta$ -mercaptoethanol  
2 $\mu$ l        LNCaP cDNA template  
5 $\mu$ l        25mM MgCl<sub>2</sub> (2.5mM final)  
30    21 $\mu$ l      dH<sub>2</sub>O  
      2 $\mu$ l        diluted Taq Polymerase (0.5U/ $\mu$ l)  
50 $\mu$ l total volume

Tubes were overlaid with 60 $\mu$ l of light mineral oil and

-49-

amplified for 30 cycles. PCR products were analyzed by electrophoresing 5 $\mu$ l of each sample on a 2-3% agarose gel followed by staining with Ethidium bromide and photography.

5    \*10x PCR Buffer

166mM NH<sub>4</sub>SO<sub>4</sub>

670mM Tris, pH 8.8

2mg/ml BSA

10   Representative photographs displaying PCR products are shown in Figure 5.

Cloning of PCR Products

15   In order to further analyze these PCR products, these products were cloned into a suitable plasmid vector using "TA Cloning" (Invitrogen® Corp.). The cloning strategy employed here is to directly ligate PCR products into a plasmid vector possessing overhanging T residues at the insertion site, exploiting the fact that Taq polymerase  
20   leaves overhanging A residues at the ends of the PCR products. The ligation mixes are transformed into competent E. coli cells and resulting colonies are grown up, plasmid DNA is isolated by the alkaline lysis method (24), and screened by restriction analysis (Figure 6 A-B).

25

DNA Sequencing of PCR Products

TA Clones of PCR products were then sequenced by the dideoxy method (25) using Sequenase (U.S. Biochemical). 3-4 $\mu$ g of each plasmid DNA was denatured with NaOH and ethanol  
30   precipitated. Labeling reactions were carried out as per the manufacturers recommendations using <sup>35</sup>S-ATP, and the reactions were terminated as per the same protocol. Sequencing products were then analyzed on 6% polyacrylamide/7M Urea gels using an IBI sequencing

- 50 -

apparatus. Gels were run at 120 watts for 2 hours. Following electrophoresis, the gels were fixed for 15-20 minutes in 10% methanol/10% acetic acid, transferred onto Whatman 3MM paper and dried down in a Biorad® vacuum dryer at 80°C for 2 hours. Gels were then autoradiographed at room temperature for 16-24 hours. In order to determine whether the PCR products were the correct clones, we analyzed the sequences obtained at the 5' and 3' ends of the molecules looking for the correct primer sequences, as well as adjacent sequences which corresponded to portions of the peptides not used in the design of the primers.

IN-20 was confirmed to be correct and represent a partial cDNA for the PSM gene. In this PCR reaction, I and N primers were used. The DNA sequence we obtained when reading from the I primer was:

ACG GAG CAA AAC TTT CAG CTT GCA AAG (SEQ ID No. 30)  
T E Q N F Q L A K (SEQ ID No. 31)

The underlined amino acids were the portion of peptide 6 that was used to design this sense primer and the remaining amino acids which agree with those present within our peptide confirm that this end of the molecule represents the correct protein (PSM antigen).

When we analyzed the other end of the molecule by reading from the N primer the sequence was:

CTC TTC GGC ATC CCA GCT TGC AAA CAA AAT TGT TCT (SEQ ID No. 32)

Since this represents the anti-sense DNA sequence, we need to show the complementary sense sequence in order to find our peptide.

-51-

Sense Sequence:

AGA ACA ATT TTG TTT GCA AGC TGG GAT GCC AAG GAG (SEQ ID No.  
33)

5     R   T   I   L   F   A   S   W   D   A   E   E (SEQ ID No.  
34)

10     The underlined amino acids here represent the portion of  
peptide 7 used to create primer N. All of the amino acids  
upstream of this primer are correct in the IN-20 clone,  
agreeing with the amino acids found in peptide 7. Further  
DNA sequencing has enabled us to identify the presence of  
our other PSM peptides within the DNA sequence of our  
positive clone.

15

The DNA sequence of this partial cDNA was found to be unique  
when screened on the Genbank computer database.

20     cDNA Library Construction and Cloning of Full - Length PSM  
cDNA

A cDNA library from LNCaP mRNA was constructed using the  
Superscript® plasmid system (BRL®-Gibco). The library was  
transformed using competent DH5-α cells and plated onto  
100mm plates containing LB plus 100μg/ml of Carbenicillin.  
25     Plates were grown overnight at 37°C and colonies were  
transferred to nitrocellulose filters. Filters were  
processed and screened as per Grunstein and Hogness (26),  
using our 1.1kb partial cDNA homologous probe which was  
radiolabelled with <sup>32</sup>P-dCTP by random priming (27). We  
30     obtained eight positive colonies which upon DNA restriction  
and sequencing analysis proved to represent full-length cDNA  
molecules coding for the PSM antigen. Shown in Figure 7 is  
an autoradiogram showing the size of the cDNA molecules  
represented in our library and in Figure 8 restriction

- 52 -

analysis of several full-length clones is shown. Figure 9 is a plasmid Southern analysis of the samples in figure 8, showing that they all hybridize to the 1.1kb partial cDNA probe.

5

Both the cDNA as well as the antigen have been screened through the Genbank Computer database (Human Genome Project) and have been found to be unique.

10 Northern Analysis of PSM Gene Expression

Northern analysis (28) of the PSM gene has revealed that expression is limited to the prostate and to prostate carcinoma.

15 RNA samples (either 10 $\mu$ g of total RNA or 2 $\mu$ g of poly A+ RNA) were denatured and electrophoresed through 1.1% agarose/formaldehyde gels at 60 milliamps for 6-8 hours. RNA was then transferred to Nytran<sup>®</sup> nylon membranes (Schleicher and Schuell<sup>®</sup>) by pressure blotting in 10x SSC  
20 with a Posi-blotter (Stratagene<sup>®</sup>). RNA was cross-linked to the membranes using a Stratalinker (Stratagene<sup>®</sup>) and subsequently baked in a vacuum oven at 80°C for 2 hours. Blots were pre-hybridized at 65°C for 2 hours in prehybridization solution (BRL<sup>®</sup>) and subsequently hybridized  
25 for 16 hours in hybridization buffer (BRL<sup>®</sup>) containing 1-2 x 10<sup>6</sup> cpm/ml of <sup>32</sup>P-labelled random-primed cDNA probe. Membranes were washed twice in 1x SSPE/1% SDS and twice in 0.1x SSPE/1% SDS at 42°C. Membranes were then air-dried and autoradiographed for 12-36 hours at -70°C.

30

PCR Analysis of PSM Gene Expression in Human Prostate Tissues

PCR was performed on 15 human prostate samples to determine PSM gene expression. Five samples each from normal prostate

-53-

tissue, benign prostatic hyperplasia, and prostate cancer were used (histology confirmed by MSKCC Pathology Department).

5 10 $\mu$ g of total RNA from each sample was reverse transcribed to made cDNA template as previously described in section IV. The primers used corresponded to the 5' and 3' ends of our 1.1kb partial cDNA, IN-20, and therefore the expected size of the amplified band is 1.1kb. Since the  $T_m$  of our primers  
10 is 64°C we annealed the primers in our PCR at 60°C. We carried out the PCR for 35 cycles using the same conditions previously described in section IV.

LNCaP and H26 - Ras transfected LNCaP (29) were included as  
15 a positive control and DU-145 as a negative control. 14/15 samples clearly amplified the 1.1kb band and therefore express the gene.

#### Experimental Results

20 The gene which encodes the 100kD PSM antigen has been identified. The complete cDNA sequence is shown in Sequence ID #1. Underneath that nucleic acid sequence is the predicted translated amino acid sequence. The total number of the amino acids is 750, ID #2. The hydrophilicity of the  
25 predicted protein sequence is shown in Figure 16. Shown in Figure 17 are three peptides with the highest point of hydrophilicity. They are: Asp-Glu-Leu-Lys-Ala-Glu (SEQ ID No. 35); Asn-Glu-Asp-Gly-Asn-Glu (SEQ ID No. 36; and Lys-Ser-Pro-Asp-Glu-Gly (SEQ ID No. 37).

30

By the method of Klein, Kanehisa and DeLisi, a specific membrane-spanning domain is identified. The sequence is from the amino acid #19 to amino acid #44: Ala-Gly-Ala-Leu-Val-Leu-Aal-Gly-Gly-Phe-Phe-Leu-Leu-Gly-Phe-Leu-Phe (SEQ ID

-54-

No. 38).

This predicted membrane-spanning domain was computed on PC Gene (computer software program). This data enables  
5 prediction of inner and outer membrane domains of the PSM antigen which aids in designing antibodies for uses in targeting and imaging prostate cancer.

When the PSM antigen sequence with other known sequences of  
10 the GeneBank were compared, homology between the PSM antigen sequence and the transferrin receptor sequence were found. The data are shown in Figure 18.

#### Experimental Discussions

15

Potential Uses for PSM Antigen:

##### 1. Tumor detection:

Microscopic:

20 Unambiguous tumor designation can be accomplished by use of probes for different antigens. For prostatic cancer, the PSM antigen probe may prove beneficial. Thus PSM could be used for diagnostic purposes and this could be accomplished at the microscopic level using in-situ hybridization using  
25 sense (control) and antisense probes derived from the coding region of the cDNA cloned by the applicants. This could be used in assessment of local extraprostatic extension, involvement of lymph node, bone or other metastatic sites. As bone metastasis presents a major problem in prostatic  
30 cancer, early detection of metastatic spread is required especially for staging. In some tumors detection of tumor cells in bone marrow portends a grim prognosis and suggests that interventions aimed at metastasis be tried. Detection of PSM antigen expression in bone marrow aspirates or  
35 sections may provide such early information. PCR



-55-

amplification or in-situ hybridization may be used. This could be developed for any possible metastatic region.

## 2. Antigenic site identification

5 The knowledge of the cDNA for the antigen also provides for the identification of areas that would serve as good antigens for the development of antibodies for use against specific amino acid sequences of the antigen. Such sequences may be at different regions such as outside,  
10-- membrane or inside of the PSM antigen. The development of these specific antibodies would provide for immunohistochemical identification of the antigen. These derived antibodies could then be developed for use, especially ones that work in paraffin fixed sections as well  
15 as frozen section as they have the greatest utility for immunodiagnosis.

## 3. Restriction fragment length polymorphism and genomic DNA

20 Restriction fragment length polymorphisms (RFLPS) have proven to be useful in documenting the progression of genetic damage that occurs during tumor initiation and promotion. It may be that RFLP analysis will demonstrate that changes in PSM sequence restriction mapping may provide  
25 evidence of predisposition to risk or malignant potential or progression of the prostatic tumor.

Depending on the chromosomal location of the PSM antigen, the PSM antigen gene may serve as a useful chromosome  
30 location marker for chromosome analysis.

## 4. Serum

With the development of antigen specific antibodies, if the antigen or selected antigen fragments appear in the serum  
35 they may provide for a serum marker for the presence of

-56-

metastatic disease and be useful individually or in combination with other prostate specific markers.

### 5. Imaging

5 As the cDNA sequence implies that the antigen has the characteristics of a membrane spanning protein with the majority of the protein on the exofacial surface, antibodies, especially monoclonal antibodies to the peptide fragments exposed and specific to the tumor may provide for  
10 tumor imaging local extension of metastatic tumor or residual tumor following prostatectomy or irradiation. The knowledge of the coding region permits the generation of monoclonal antibodies and these can be used in combination to provide for maximal imaging purposes. Because the  
15 antigen shares a similarity with the transferrin receptor based on cDNA analysis (approximately 54%), it may be that there is a specific normal ligand for this antigen and that identification of the ligand(s) would provide another means of imaging.

20

### 6. Isolation of ligands

The PSM antigen can be used to isolate the normal ligand(s) that bind to it. These ligand(s) depending on specificity may be used for targeting, or their serum levels may be  
25 predictive of disease status. If it is found that the normal ligand for PSM is a carrier molecule then it may be that PSM could be used to bind to that ligand for therapy purposes (like an iron chelating substance) to help remove the ligand from the circulation. If the ligand promotes  
30 tumor growth or metastasis then providing soluble PSM antigen would remove the ligand from binding the prostate. Knowledge of PSM antigen structure could lend to generation of small fragment that binds ligand which could serve the same purpose.

35

-57-

## 7. Therapeutic uses

a) Ligands. The knowledge that the cDNA structure of PSM antigen shares structural homology with the transferrin receptor (54% on the nucleic acid level) implies that there may be an endogenous ligand for the receptor that may or may not be transferrin-like. Transferrin is thought to be a ligand that transports iron into the cell after binding to the transferrin receptor. However, apotransferrin is being reported to be a growth factor for some cells which express the transferrin receptor (30). Whether transferrin is a ligand for this antigen or some other ligand binds to this ligand remains to be determined. If a ligand is identified it may carry a specific substance such as a metal ion (iron or zinc or other) into the tumor and thus serve as a means to deliver toxic substances (radioactive or cytotoxic chemical i.e. toxin like ricin or cytotoxic alkylating agent or cytotoxic prodrug) to the tumor.

The main metastatic site for prostatic tumor is the bone. The bone and bone stroma are rich in transferrin. Recent studies suggest that this microenvironment is what provides the right "soil" for prostatic metastasis in the bone (31). It may be that this also promotes attachment as well, these factors which reduce this ability may diminish prostatic metastasis to the bone and prostatic metastatic growth in the bone.

It was found that the ligand for the new antigen (thought to be an oncogene and marker of malignant phenotype in breast carcinoma) served to induce differentiation of breast cancer cells and thus could serve as a treatment for rather than promotor of the disease. It may be that ligand binding to the right region of PSM whether with natural ligand or with an antibody may serve a similar function.

- 58 -

- Antibodies against PSM antigen coupled with a cytotoxic agent will be useful to eliminate prostate cancer cells. Transferrin receptor antibodies with toxin conjugates are cytotoxic to a number of tumor cells as tumor cells tend to express increased levels of transferrin receptor (32). Transferrin receptors take up molecules into the cell by endocytosis. Antibody drug combinations can be toxic. Transferrin linked toxin can be toxic.
- 10 \_ b) Antibodies against PSM antigen coupled with a cytotoxic agent will be useful to eliminate prostate cancer cells. The cytotoxic agent may be a radioisotope or toxin as known in ordinary skill of the art. The linkage of the antibody and the toxin or radioisotope can be chemical. Examples of
- 15 direct linked toxins are doxorubicin, chlorambucil, ricin, pseudomonas exotoxin etc., or a hybrid toxin can be generated  $\frac{1}{2}$  with specificity for PSM and the other  $\frac{1}{2}$  with specificity for the toxin. Such a bivalent molecule can serve to bind to the tumor and the other  $\frac{1}{2}$  to deliver a
- 20 cytotoxic to the tumor or to bind to and activate a cytotoxic lymphocyte such as binding to the  $T_1$  -  $T_3$  receptor complex. Antibodies of required specificity can also be cloned into T cells and by replacing the immunoglobulin domain of the T cell receptor (TCR); cloning in the desired
- 25 MAb heavy and light chains; splicing the  $U_h$  and  $U_l$  gene segments with the constant regions of the  $\alpha$  and  $\beta$  TCR chains and transfecting these chimeric Ab/TCR genes in the patients' T cells, propagating these hybrid cells and infusing them into the patient (33). Specific knowledge of
- 30 tissue specific antigens for targets and generation of MAb's specific for such targets will help make this a usable approach. Because the PSM antigen coding region provides knowledge of the entire coding region, it is possible to generate a number of antibodies which could then be used in

-59-

combination to achieve an additive or synergistic anti-tumor action. The antibodies can be linked to enzymes which can activate non-toxic prodrugs at its site of the tumor such as Ab-carboxypeptidase and 4-(bis(2 chloroethyl)amino)benzoyl-  
5  $\alpha$ -glutamic acid and its active parent drug in mice (34).

It is possible to produce a toxic genetic chimera such as TP-40 a genetic recombinant that possesses the cDNA from TGF- $\alpha$  and the toxic portion of pseudomonas exotoxin so  
10 the TGF and portion of the hybrid binds the epidermal growth factor receptor (EGFR) and the pseudomonas portion gets taken up into the cell enzymatically and inactivates the ribosomes ability to perform protein synthesis resulting in cell death. When we know the ligand for the PSM antigen we  
15 can do the same.

In addition, once the ligand for the PSM antigen is identified, toxin can be chemically conjugated to the ligands. Such conjugated ligands can be therapeutically  
20 useful. Examples of the toxins are daunomycin, chlorambucil, ricin, pseudomonas exotoxin, etc. Alternatively, chimeric construct can be created linking the cDNA of the ligand with the cDNA of the toxin. An example of such toxin is TGF $\alpha$  and pseudomonas exotoxin (35).

25

#### 8. Others

The PSM antigen may have other uses. It is well known that the prostate is rich in zinc, if the antigen provides function relative to this or other biologic function the PSM  
30 antigen may provide for utility in the treatment of other prostatic pathologies such as benign hyperplastic growth and/or prostatitis.

Because purified PSM antigen can be generated, the purified  
35 PSM antigen can be linked to beads and use it like a

-60-

standard "affinity" purification. Serum, urine or other biological samples can be used to incubate with the PSM antigen bound onto beads. The beads may be washed thoroughly and then eluted with salt or pH gradient. The  
5 eluted material is SDS gel purified and used as a sample for microsequencing. The sequences will be compared with other known proteins and if unique, the technique of degenerated PCR can be employed for obtaining the ligand. Once known, the affinity of the ligand will be determined by standard  
10-- protocols (15).

References of the First Series of Experiments

1. Chiaroda, A. (1991) National roundtable of prostate cancer: research directions. Cancer Res. 51: 2498-2505.  
5
2. Coffey, D.S. Prostate Cancer - An overview of an increasing dilemma. Cancer Supplement, 71,3: 880-886, 1993.  
10
3. Warner, J.A., et al., (1991) Future developments of non-hormonal systemic therapy for prostatic carcinoma. Urologic Clin. North Amer. 18:25-33.
- 15 4. Nguyen, L., et al., (1990) Prostatic acid phosphatase in the serum of cancer patients with prostatic cancer is a specific phosphotyrosine acid phosphatase. Clin. Chem. 35:1450-1455.
- 20 5. Henttu, P., et al., (1989) cDNA coding for the entire human prostate specific antigen show high homologies to the human tissue kallikrein genes. Bioch. Biophys. Res. Comm. 160:903-908.
- 25 6. Yong, CY-F., et al., (1991) Hormonal regulation of prostate-specific antigen messenger RNA in human prostatic adenocarcinoma cell line LNCaP. Cancer Res. 51:3748-3752.
- 30 7. Liotta, L.A. (1986) Tumor invasion and metastases: role of the extracellular matrix. Cancer Res. 46:1-7.
8. Horoszewicz, J.S., et al. (1987) Monoclonal antibodies to a new antigenic marker in epithelial prostatic cells and serum of prostatic cancer patients. Anticancer  
35

-62-

Res. 7:927-936.

9. Horoszewicz, J.S., et al. (1983) LNCaP model of human prostatic carcinoma. Cancer Res., 43:1809-1818.
- 5 10. Lopes, D., et al. (1990) Immunohistochemical and pharmacokinetic characterization of the site-specific immunoconjugate CYT-356, derived from anti-prostate monoclonal antibody 7E11-C5. Cancer Res., 50:6423-10 6429.
11. Wright, Jr., et al., (1990) Characterization of a new carcinoma associated marker:7E11-C5. Antibod. Immunconj. Radiopharm.3:(abst#193).
- 15 12. Feng, Q., et al., (1991) Purification and biochemical characterization of the 7E11-C5 prostate carcinoma associated antigen. Proc. Amer. Assoc. Cancer Res. 32:239.
- 20 13. Axelrod, H.R., et al., Preclinical results and human immunohistochemical studies with <sup>90</sup>Y-CYT-356. A New prostate cancer agent. Abstract 596. AUA 87th Annual Meeting, May 10-14, 1992. Washington, D.C.
- 25 14. Maniatis, T., et al., (1982) Molecular Cloning; Cold Spring Harbor Laboratory, pp.197-98 (1982).
- 30 15. Maniatis, et al., (1982) Molecular Cloning, Cold Spring Harbor Laboratory.
16. Methods in Enzymology vol. 34: 1-810, 1974 (E) B. Jacoby and M. Wilchek Academic Press, New York 1974.



-63-

17. Hogan B. et al. (1986) Manipulating the Mouse Embryo, A Laboratory Manual, Cold Spring Harbor Laboratory.
18. Capecchi M.R. Science (1989) 244:1288-1292; Zimmer, A.  
5 and Gruss, P. (1989) Nature 338:150-153.
19. Trowbridge, I.S., (1982) Prospects for the clinical use of cytotoxic monoclonal antibodies conjugates in the treatment of cancer. Cancer Surveys 1:543-556.
10. 20. Hank, S.K. (1987) Homology probing: Identification of cDNA clones encoding members of the protein-serine kinase family. Proc. Natl. Acad. Sci. 84:388-392.
- 15 21. Lee, C.C., et al., (1988) Generation of cDNA probes directed by amino acid sequences: cloning of urate oxidase. Science, 239, 1288.
- 20 22. Girgis, S.I., et al. (1988) Generation of DNA probes for peptides with highly degenerate codons using mixed primer PCR. Nucleic Acids Res. 16:10932.
- 25 23. Kartner, N., et al. (1977) Isolation of plasma membranes from human skin fibroblasts. J. Membrane Biology, 36:191-211.
- 30 24. Hsu, S.M., et al. (1981) Comparative study of the immunoperoxidase, anti-peroxidase, and avidin-biotin-complex method for studying polypeptide hormones with radioimmunoassay antibodies. Am. J. Pathology, 75:734.
- 35 25. Tempst, P., et al. (1989) Examination of automated polypeptide sequencing using standard phenylisothiocyanate reagent and subpicomole high performance liquid chromatography analysis. Analytical

-64-

Biochem. 183:290-300.

26. Birnboim, H.C. (1983) A rapid alkaline extraction  
method for the isolation of plasmid DNA. Meth.  
5 Enzymol, 100:243-255.
27. Sanger, F., et al. (1977) DNA sequencing with chain-  
terminating inhibitors. Proc. Natl. Acad. Sci. USA,  
74:5463-5467.
- 10 28. Grunstein, M., et al. (1975) Colony hybridization as a  
method for the isolation of cloned DNAs that contain a  
specific gene. Proc. Natl. Acad. Sci. USA, 72:3961.
- 15 29. Feinberg, A.P., et al. (1983) A technique for  
radiolabeling DNA restriction endonuclease fragments to  
high specific activity. Anal. Biochem, 132, 6.
- 20 30. Rave, N., et al. (1979) Identification of procollagen  
mRNAs transferred to diazobenzylomethyl paper from  
formaldehyde gels. Nucleic Acids Research, 6:3559.
- 25 31. Voeller, H.J., et al. (1991) v-ras<sup>H</sup> expression confers  
hormone-independent in-vitro growth to LNCaP prostate  
carcinoma cells. Molec. Endocrinology. Vol. 5. No. 2,  
209-216.
- 30 32. Sirbasku, D.A. (1991) Purification of an equine  
apotransferrin variant (thyromedin) essential for  
thyroid hormone dependent growth of GH<sub>1</sub>, rat pituitary  
tumor cells in chemically defined culture. Biochem.,  
30:295-301.
33. Rossi, M.C. (1992) Selective stimulation of prostatic

-65-

carcinoma cell proliferation by transferrin. Proc. Natl. Acad. Sci. (USA) 89:6197-6201.

- 5 34. Eshhan, Z. (1990) Chimeric T cell receptor which incorporates the anti-tumor specificity of a monoclonal antibody with the cytolytic activity of T cells: a model system for immunotherapeutic approach. B. J. Cancer 62:27-29.
- 10 35. Antonie, P. (1990) Disposition of the prodrug 4-(bis(2 chloroethyl) amino)benzoyl- $\alpha$ -glutamic acid and its active parent in mice. B. J. Cancer 62:905-914.
- 15 36. Heimbrook, D.C., et al. (1990) Transforming growth factor alpha-pseudomonas exotoxin fusion protein prolongs survival of nude mice bearing tumor xenografts. Proc. Natl. Acad. Sci. (USA) 87:4697-4701.
- 20 37. Chiarodo, A. National Cancer Institute roundtable on prostate cancer; future research directions. Cancer Res., 51: 2498-2505, 1991.
- 25 38. Abdel-Nabi, H., Wright, G.L., Gulfo, J.V., Petrylak, D.P., Neal, C.E., Texter, J.E., Begun, F.P., Tyson, I., Heal, A., Mitchell, E., Purnell, G., and Harwood, S.J. Monoclonal antibodies and radioimmunoconjugates in the diagnosis and treatment of prostate cancer. Semin. Urol., 10: 45-54, 1992.

- 66 -

Second Series of ExperimentsExpression of the Prostate-Specific Membrane Antigen

5 Applicant's have recently cloned a 2.65 kb complementary DNA encoding PSM, the prostate-specific membrane antigen recognized by the 7E11-C5.3 anti-prostate monoclonal antibody. Immunohistochemical analysis of the LNCaP, DU-145, and PC-3 prostate cancer cell lines for PSM expression  
10 using the 7E11-C5.3 antibody reveals intense staining in the LNCaP cells, with no detectable expression in both the DU-145 and PC-3 cells. Coupled in-vitro transcription/translation of the 2.65 kb full-length PSM cDNA yields an 84 kDa protein corresponding to the predicted polypeptide  
15 molecular weight of PSM. Post-translational modification of this protein with pancreatic canine microsomes yields the expected 100 kDa PSM antigen. Following transfection of PC-3 cells with the full-length PSM cDNA in a eukaryotic expression vector applicant's detect expression of the PSM  
20 glycoprotein by Western analysis using the 7E11-C5.3 monoclonal antibody. Ribonuclease protection analysis demonstrates that the expression of PSM mRNA is almost entirely prostate-specific in human tissues. PSM expression appears to be highest in hormone-deprived states and is  
25 hormonally modulated by steroids, with DHT downregulating PSM expression in the human prostate cancer cell line LNCaP by 8-10 fold, testosterone downregulating PSM by 3-4 fold, and corticosteroids showing no significant effect. Normal and malignant prostatic tissues consistently show high PSM  
30 expression, whereas we have noted heterogeneous, and at times absent, expression of PSM in benign prostatic hyperplasia. LNCaP tumors implanted and grown both orthotopically and subcutaneously in nude mice, abundantly express PSM providing an excellent in-vivo model system to  
35 study the regulation and modulation of PSM expression.

## Experimental Details

### Materials and Methods

#### 5 Cells and Reagents:

The LNCaP, DU-145, and PC-3 cell lines were obtained from the American Type Culture Collection. Details regarding the establishment and characteristics of these cell lines have been previously published (5A,7A,8A). Unless specified  
10 otherwise, LNCaP cells were grown in RPMI 1640 media supplemented with L-glutamine, nonessential amino acids, and 5% fetal calf serum (Gibco-BRL, Gaithersburg, MD.) in a CO<sub>2</sub> incubator at 37C. DU-145 and PC-3 cells were grown in  
15 minimal essential medium supplemented with 10% fetal calf serum. All cell media were obtained from the MSKCC Media Preparation Facility. Restriction and modifying enzymes were purchased from Gibco-BRL unless otherwise specified.

#### Immunohistochemical Detection of PSM

20 We employed the avidin-biotin method of detection to analyze prostate cancer cell lines for PSM antigen expression (9A). Cell cytopspins were made on glass slides using  $5 \times 10^4$  cells/100ul per slide. Slides were washed twice with PBS and then incubated with the appropriate suppressor serum for  
25 20 minutes. The suppressor serum was drained off and the cells were incubated with diluted 7E11-C5.3 (5g/ml) monoclonal antibody for 1 hour. Samples were then washed with PBS and sequentially incubated with secondary  
30 antibodies for 30 minutes and with avidin-biotin complexes for 30 minutes. Diaminobenzidine served as our chromogen and color development followed by hematoxylin counterstaining and mounting. Duplicate cell cytopspins were used as controls for each experiment. As a positive control, the anti-cytokeratin monoclonal antibody CAM 5.2

-68-

was used following the same procedure described above. Human EJ bladder carcinoma cells served as a negative control.

5    In-Vitro Transcription/Translation of PSM Antigen

Plasmid 55A containing the full length 2.65 kb PSM cDNA in the plasmid pSPORT 1 (Gibco-BRL) was transcribed in-vitro using the Promega TNT system (Promega Corp. Madison, WI). T7 RNA polymerase was added to the cDNA in a reaction mixture containing rabbit reticulocyte lysate, an amino acid mixture lacking methionine, buffer, and <sup>35</sup>S-Methionine (Amersham) and incubated at 30C for 90 minutes. Post-translational modification of the resulting protein was accomplished by the addition of pancreatic canine microsomes into the reaction mixture (Promega Corp. Madison, WI.). Protein products were analyzed by electrophoresis on 10% SDS-PAGE gels which were subsequently treated with Amplify autoradiography enhancer (Amersham, Arlington Heights, IL.) according to the manufacturers instructions and dried at 80C in a vacuum dryer. Gels were autoradiographed overnight at -70C using Hyperfilm MP (Amersham).

Transfection of PSM into PC-3 Cells

The full length PSM cDNA was subcloned into the pREP7 eukaryotic expression vector (Invitrogen, San Diego, CA.). Plasmid DNA was purified from transformed DH5-alpha bacteria (Gibco-BRL) using Qiagen maxi-prep plasmid isolation columns (Qiagen Inc., Chatsworth, CA.). Purified plasmid DNA (6-10g) was diluted with 900ul of Optimem media (Gibco-BRL) and mixed with 30ul of Lipofectin reagent (Gibco-BRL) which had been previously diluted with 900ul of Optimem media. This mixture was added to T-75 flasks of 40-50% confluent PC-3 cells in Optimem media. After 24-36 hours, cells were trypsinized and split into 100mm dishes containing RPMI

-69-

1640 media supplemented with 10% fetal calf serum and 1 mg/ml of Hygromycin B (Calbiochem, La Jolla, CA.). The dose of Hygromycin B used was previously determined by a time course/dose response cytotoxicity assay. Cells were maintained in this media for 2-3 weeks with changes of media and Hygromycin B every 4-5 days until discrete colonies appeared. Colonies were isolated using 6mm cloning cylinders and expanded in the same media. As a control, PC-3 cells were also transfected with the pREP7 plasmid alone. RNA was isolated from the transfected cells and PSM mRNA expression was detected by both RNase Protection analysis (described later) and by Northern analysis.

#### Western Blot Detection of PSM Expression

Crude protein lysates were isolated from LNCaP, PC-3, and PSM-transfected PC-3 cells as previously described (10A). LNCaP cell membranes were also isolated according to published methods (10A). Protein concentrations were quantitated by the Bradford method using the BioRad protein reagent kit (BioRad, Richmond, CA.). Following denaturation, 20g of protein was electrophoresed on a 10% SDS-PAGE gel at 25 mA for 4 hours. Gels were electroblotted onto Immobilon P membranes (Millipore, Bedford, MA.) overnight at 4C. Membranes were blocked in 0.15M NaCl/0.01M Tris-HCl (TS) plus 5% BSA followed by a 1 hour incubation with 7E11-C5.3 monoclonal antibody (10g/ml). Blots were washed 4 times with 0.15M NaCl/0.01M Tris-HCl/0.05% Triton-X 100 (TS-X) and incubated for 1 hour with rabbit anti-mouse IgG (Accurate Scientific, Westbury, N.Y.) at a concentration of 10g/ml.

Blots were then washed 4 times with TS-X and labeled with <sup>125</sup>I-Protein A (Amersham, Arlington Heights, IL.) at a concentration of 1 million cpm/ml. Blots were then washed 4 times with TS-X and dried on Whatman 3MM paper, followed

-70-

by overnight autoradiography at -70C using Hyperfilm MP (Amersham).

5    Orthotopic and Subcutaneous LNCaP Tumor Growth in Nude Mice  
LNCaP cells were harvested from sub-confluent cultures by a  
one minute exposure to a solution of 0.25% trypsin and 0.02%  
EDTA. Cells were resuspended in RPMI 1640 media with 5%  
fetal bovine serum, washed and diluted in either Matrigel  
10    (Collaborative Biomedical Products, Bedford, MA.) or calcium  
and magnesium-free Hank's balanced salt solution (HBSS).  
Only single cell suspensions with greater than 90% viability  
by trypan blue exclusion were used for in vivo injection.  
Male athymic Swiss (nu/nu) nude mice 4-6 weeks of age were  
15    obtained from the Memorial Sloan-Kettering Cancer Center  
Animal Facility. For subcutaneous tumor cell injection one  
million LNCaP cells resuspended in 0.2 mls. of Matrigel were  
injected into the hindlimb of each mouse using a disposable  
syringe fitted with a 28 gauge needle. For orthotopic  
20    injection, mice were first anesthetized with an  
intraperitoneal injection of Pentobarbital and placed in the  
supine position. The abdomen was cleansed with Betadine and  
the prostate was exposed through a midline incision. 2.5  
million LNCaP tumor cells in 0.1 ml. were injected directly  
25    into either posterior lobe using a 1 ml disposable syringe  
and a 28 gauge needle. LNCaP cells with and without  
Matrigel were injected. Abdominal closure was achieved in  
one layer using Autoclip wound clips (Clay Adams,  
Parsippany, N.J.). Tumors were harvested in 6-8 weeks,  
30    confirmed histologically by faculty of the Memorial Sloan-  
Kettering Cancer Center Pathology Department, and frozen in  
liquid nitrogen for subsequent RNA isolation.

#### RNA Isolation

35    Total cellular RNA was isolated from cells and tissues by  
standard techniques (11,12) as well as by using RNazol B



- 71 -

(Cinna/Biotechx, Houston, TX.). RNA concentrations and quality were assessed by UV spectroscopy on a Beckman DU 640 spectrophotometer and by gel analysis. Human tissue total RNA samples were purchased from Clontech Laboratories, Inc.,  
5 Palo Alto, CA.

#### Ribonuclease Protection Assays

A portion of the PSM cDNA was subcloned into the plasmid vector pSPORT 1 (Gibco-BRL) and the orientation of the cDNA  
10 insert relative to the flanking T7 and SP6 RNA polymerase promoters was verified by restriction analysis. Linearization of this plasmid upstream of the PSM insert followed by transcription with SP6 RNA polymerase yields a 400 nucleotide antisense RNA probe, of which 350 nucleotides  
15 should be protected from RNase digestion by PSM RNA. This probe was used in figure 20. Plasmid IN-20, containing a 1 kb partial PSM cDNA in the plasmid pCR II (Invitrogen) was also used for riboprobe synthesis. IN-20 linearized with Xmn I (Gibco-BRL) yields a 298 nucleotide anti-sense RNA  
20 probe when transcribed using SP6 RNA polymerase, of which 260 nucleotides should be protected from RNase digestion by PSM mRNA. This probe was used in figures 21 and 22. Probes were synthesized using SP6 RNA polymerase (Gibco-BRL), rNTPs (Gibco-BRL), RNasin (Promega), and <sup>32</sup>P-rCTP (NEN, Wilmington, DE.) according to published protocols (13). Probes were  
25 purified over NENSORB 20 purification columns (NEN) and approximately 1 million cpm of purified, radiolabeled PSM probe was mixed with 10g of each RNA and hybridized overnight at 45C using buffers and reagents from the RPA II  
30 kit (Ambion, Austin, TX). Samples were processed as per manufacturer's instructions and analyzed on 5% polyacrilamide/7M urea denaturing gels using Seq ACRYL reagents (ISS, Natick, MA.). Gels were pre-heated to 55C and run for approximately 1-2 hours at 25 watts. Gels were

- 72 -

then fixed for 30 minutes in 10% methanol/10% acetic acid, dried onto Whatman 3MM paper at 80C in a BioRad vacuum dryer and autoradiographed overnight with Hyperfilm MP (Amersham). Quantitation of PSM expression was determined by using a scanning laser densitometer (LKB, Piscataway, NJ.).

#### Steroid Modulation Experiment

LNCaP cells (2 million) were plated onto T-75 flasks in RPMI 1640 media supplemented with 5% fetal calf serum and grown 24 hours until approximately 30-40% confluent. Flasks were then washed several times with phosphate-buffered saline and RPMI medium supplemented with 5% charcoal-extracted serum was added. Cells were then grown for another 24 hours, at which time dihydrotestosterone, testosterone, estradiol, progesterone, and dexamethasone (Steraloids Inc., Wilton, NH.) were added at a final concentration of 2 nM. Cells were grown for another 24 hours and RNA was then harvested as previously described and PSM expression analyzed by ribonuclease protection analysis.

#### Experimental Results

##### **Immunohistochemical Detection of PSM:**

Using the 7E11-C5.3 anti-PSM monoclonal antibody, PSM expression is clearly detectable in the LNCaP prostate cancer cell line, but not in the PC-3 and DU-145 cell lines (figure 17) in agreement with previously published results (4A). All normal and malignant prostatic tissues analyzed stained positively for PSM expression (unpublished data).

##### **In-Vitro Transcription/Translation of PSM Antigen:**

As shown in figure 18, coupled in-vitro transcription/translation of the 2.65 kb full-length PSM cDNA yields an 84 kDa protein species in agreement with the expected protein

-73-

product from the 750 amino acid PSM open reading frame. Following post-translational modification using pancreatic canine microsomes we obtained a 100 kDa glycosylated protein species consistent with the mature, native PSM antigen.

#### Detection of PSM Antigen in LNCaP Cell Membranes and Transfected PC-3 Cells:

PC-3 cells transfected with the full length PSM cDNA in the pREP7 expression vector were assayed for expression of SM mRNA by Northern analysis (data not shown). A clone with high PSM mRNA expression was selected for PSM antigen analysis by Western blotting using the 7E11-C5.3 antibody. In figure 19, the 100 kDa PSM antigen is well expressed in LNCaP cell lysate and membrane fractions, as well as in PSM-transfected PC-3 cells but not in native PC-3 cells. This detectable expression in the transfected PC-3 cells proves that the previously cloned 2.65 kb PSM cDNA encodes the antigen recognized by the 7E11-C5.3 anti-prostate monoclonal antibody and that the antigen is being appropriately glycosylated in the PC-3 cells, since the antibody recognizes a carbohydrate-containing epitope on PSM.

#### PSM mRNA Expression:

Expression of PSM mRNA in normal human tissues was analyzed using ribonuclease protection assays. Tissue expression of PSM appears predominantly within the prostate, with very low levels of expression detectable in human brain and salivary gland (figure 20). No detectable PSM mRNA expression was evident in non-prostatic human tissues when analyzed by Northern analysis (data not shown). We have also noted on occasion detectable PSM expression in normal human small intestine tissue, however this mRNA expression is variable depending upon the specific riboprobe used (data not shown). All samples of normal human prostate and human prostatic

- 74 -

adenocarcinoma assayed have revealed clearly detectable PSM expression, whereas we have noted generally decreased or absent expression of PSM in tissues exhibiting benign hyperplasia (figure 21). In human LNCaP tumors grown both  
5 orthotopically and subcutaneously in nude mice we detected abundant PSM expression with or without the use of matrigel, which is required for the growth of subcutaneously implanted LNCaP cells (figure 21). PSM mRNA expression is distinctly modulated by the presence of steroids in physiologic doses  
10 (figure 22). DHT downregulated expression by 8-10 fold after 24 hours and testosterone diminished PSM expression by 3-4 fold. Estradiol and progesterone also downregulated PSM expression in LNCaP cells, perhaps as a result of binding to the mutated androgen receptor known to exist in the LNCaP  
15 cell. Overall, PSM expression is highest in the untreated LNCaP cells grown in steroid-depleted media, a situation that we propose simulates the hormone-deprived (castrate) state in-vivo. This experiment was repeated at steroid dosages ranging from 2-200 nM and at time points from 6  
20 hours to 7 days with similar results; maximal downregulation of PSM mRNA was seen with DHT at 24 hours at doses of 2-20 nM.

#### Experimental Discussion

25 In order to better understand the biology of the human prostate in both normal and neoplastic states, we need to enhance our knowledge by studying the various proteins and other features that are unique to this important gland. Previous research has provided two valuable prostatic bio-  
30 markers, PAP and PSA, both of which have had a significant impact on the diagnosis, treatment, and management of prostate malignancies. Our present work describing the preliminary characterization of the prostate-specific membrane antigen (PSM) reveals it to be a gene with many  
35 interesting features. PSM is almost entirely prostate-

-75-

specific as are PAP and PSA, and as such may enable further delineation of the unique functions and behavior of the prostate. The predicted sequence of the PSM protein (3) and its presence in the LNCaP cell membrane as determined by Western blotting and immunohistochemistry, indicate that it is an integral membrane protein. Thus, PSM provides an attractive cell surface epitope for antibody-directed diagnostic imaging and cytotoxic targeting modalities (14). The ability to synthesize the PSM antigen *in-vitro* and to produce tumor xenografts maintaining high levels of PSM expression provides us with a convenient and attractive model system to further study and characterize the regulation and modulation of PSM expression. Also, the high level of PSM expression in the LNCaP cells provides an excellent *in-vitro* model system. Since PSM expression is hormonally-responsive to steroids, and may be highly expressed in hormone-refractory disease (15), it is imperative to elucidate the potential role of PSM in the evolution of androgen-independent prostate cancer. The detection of PSM mRNA expression in minute quantities in brain, salivary gland, and small intestine warrants further investigation, although these tissues were negative for expression of PSM antigen by immunohistochemistry using the 7E11-C5.3 antibody (16). In all of these tissues, particularly small intestine, we detected mRNA expression using a probe corresponding to a region of the PSM cDNA near the 3' end, whereas we were unable to detect expression when using a 5' end PSM probe. These results may indicate that the PSM mRNA transcript undergoes alternative splicing in different tissues. Previous protein studies have suggested that the 7E11-C5.3 antibody may actually detect two other slightly larger protein species in addition to the 100 kDa PSM antigen (17). These other protein species can be seen in the LNCaP lysate and membrane samples in figure 19. Possible origins of these proteins include alternatively

-76-

spliced PSM mRNA, other genes distinct from but closely related to PSM, or different post-translational modifications of the PSM protein. We are currently investigating these possibilities.

5

Applicnat's approach is based on prostate tissue specific promotor:enzyme or cytokine chimeras. We will examine promotor specific activation of prodrugs such as non toxic gancyclovir which is converted to a toxic metabolite by  
10 herpes simplex thymidine kinase or the prodrug 4-(bis(2chloroethyl)amino)benzoyl-1-glutamic acid to the benzoic acid mustard alkylating agent by the pseudomonas carboxy peptidase G2. As these drugs are activated by the enzyme (chimera) specifically in the tumor the active drug  
15 is released only locally in the tumor environment, destroying the surrounding tumor cells. We will also examine the promotor specific activation of cytokines such as IL-12, IL-2 or GM-CSF for activation and specific antitumor vaccination. Lastly the tissue specific promotor  
20 activation of cellular death genes may also prove to be useful in this area.

#### Gene Therapy Chimeras

The establishment of "chimeric DNA" for gene therapy  
25 requires the joining of different segments of DNA together to make a new DNA that has characteristics of both precursor DNA species involved in the linkage. In this proposal the two pieces being linked involve different functional aspects of DNA, the promotor region which allows for the reading of  
30 the DNA for the formation of mRNA will provide specificity and the DNA sequence coding for the mRNA will provide for therapeutic functional DNA.

#### DNA-Specified Enzyme or Cytokine mRNA:

35 When effective, antitumor drugs can cause the regression of

- 77 -

very large amounts of tumor. The main requirements for antitumor drug activity is the requirement to achieve both a long enough time (t) and high enough concentration (c) (cxt) of exposure of the tumor to the toxic drug to assure  
5 sufficient cell damage for cell death to occur. The drug also must be "active" and the toxicity for the tumor greater than for the hosts normal cells (22). The availability of the drug to the tumor depends on tumor blood flow and the drugs diffusion ability. Blood flow to the tumor does not  
10 provide for selectivity as blood flow to many normal tissues is often as great or greater than that to the tumor. The majority of chemotherapeutic cytotoxic drugs are often as toxic to normal tissue as to tumor tissue. Dividing cells are often more sensitive than non-dividing normal cells, but  
15 in many slow growing solid tumors such as prostatic cancer this does not provide for antitumor specificity (22).

Previously a means to increase tumor specificity of antitumor drugs was to utilize tumor associated enzymes to  
20 activate nontoxic prodrugs to cytotoxic agents (19). A problem with this approach was that most of the enzymes found in tumors were not totally specific in their activity and similar substrate active enzymes or the same enzyme at only slightly lower amounts was found in other tissue and  
25 thus normal tissues were still at risk for damage.

To provide absolute specificity and unique activity, viral, bacterial and fungal enzymes which have unique specificity for selected prodrugs were found which were not present in  
30 human or other animal cells. Attempts to utilize enzymes such as herpes simplex thymidine kinase, bacterial cytosine deaminase and carboxypeptidase G-2 were linked to antibody targeting systems with modest success (19). Unfortunately, antibody targeted enzymes limit the number of enzymes  
35 available per cell. Also, most antibodies do not have a

- 78 -

high tumor target to normal tissue ratio thus normal tissues are still exposed reducing the specificity of these unique enzymes. Antibodies are large molecules that have poor diffusion properties and the addition of the enzymes  
5 molecular weight further reduces the antibodies diffusion.

Gene therapy could produce the best desired result if it could achieve the specific expression of a protein in the tumor and not normal tissue in order that a high local  
10 concentration of the enzyme be available for the production in the tumor environment of active drug (21).

#### Cytokines:

Applicant's research group has demonstrated that Applicant's  
15 can specifically and non-toxically "cure" an animal of an established tumor, in models of bladder or prostate cancer. The prostate cancer was the more difficult to cure especially if it was grown orthotopically in the prostate.

20 Our work demonstrated that tumors such as the bladder and prostate were not immunogenic, that is the administration of irradiated tumor cells to the animal prior to subsequent administration of non-irradiated tumor cells did not result in a reduction of either the number of tumor cells to  
25 produce a tumor nor did it reduce the growth rate of the tumor. But if the tumor was transfected with a retrovirus and secreted large concentrations of cytokines such as IL-2 then this could act as an antitumor vaccine and could also reduce the growth potential of an already established and  
30 growing tumor. IL-2 was the best, GM-CSF also had activity whereas a number of other cytokines were much less active. In clinical studies just using IL-2 for immunostimulation, very large concentrations had to be given which proved to be toxic. The key to the success of the cytokine gene modified  
35 tumor cell is that the cytokine is produced at the tumor



- 79 -

site locally and is not toxic and that it stimulates immune recognition of the tumor and allows specific and non toxic recognition and destruction of the tumor. The exact mechanisms of how IL-2 production by the tumor cell  
5 activates immune recognition is not fully understood, but one explanation is that it bypasses the need for cytokine production by helper T cells and directly stimulates tumor antigen activated cytotoxic CD8 cells. Activation of antigen presenting cells may also occur.

10 --

#### Tissue Promotor-Specific Chimera DNA Activation

##### **Non-Prostatic Tumor Systems:**

It has been observed in non-prostatic tumors that the use of  
15 promotor specific activation can selectively lead to tissue specific gene expression of the transfected gene. In melanoma the use of the tyrosinase promotor which codes for the enzyme responsible for melanin expression produced over a 50 fold greater expression of the promotor driven reporter  
20 gene expression in melanoma cells and not non melanoma cells. Similar specific activation was seen in the melanoma cells transfected when they were growing in mice. In that experiment no non-melanoma or melanocyte cell expressed the tyrosinase drive reporter gene product. The research group  
25 at Wellcome Laboratories have cloned and sequenced the promoter region of the gene coding for carcinoembryonic antigen (CEA). CEA is expressed on colon and colon carcinoma cells but specifically on metastatic cytosine deaminase which converts 5 flurorocytosine into 5  
30 fluorouracil and observed a large increase in the ability to selectively kill CEA promotor driven colon tumor cells but non dividing not dividing normal liver cells. In vivo they observed that bystander tumor cells which were not transfected with the cytosine deaminase gene were also  
35 killed, and that there was no toxicity to the host animal as

-80-

the large tumors were regressing following treatment. Herpes simplex virus, (HSV), thymidine kinase similarly activates the prodrug gancyclovir to be toxic towards dividing cancer cells and HSV thymidine kinase has been  
5 shown to be specifically activatable by tissue specific promoters.

#### Prostatic Tumor Systems:

The therapeutic key to effective cancer therapy is to  
10 achieve specificity and spare the patient toxicity. Gene therapy may provide a key part to specificity in that non-essential tissues such as the prostate and prostatic tumors produce tissue specific proteins, such as acid phosphatase (PAP), prostate specific antigen (PSA), and a gene which we  
15 cloned, prostate-specific membrane antigen (PSM). Tissues such as the prostate contain selected tissue specific transcription factors which are responsible for binding to the promoter region of the DNA of these tissue specific mRNA. The promoter for PSA has been cloned and we are  
20 investigating its use as a prostate specific promotor for prostatic tumor cells. Usually patients who are being treated for metastatic prostatic cancer have been put on androgen deprivation therapy which dramatically reduces the expression of mRNA for PSA. PSM on the other hand increases  
25 in expression with hormone deprivation which-means it would be even more intensely expressed on patients being treated with hormone therapy. Preliminary work in collaboration with Dr. John Isaacs' Laboratory demonstrates that PSM is expressed when the human chromosome region containing the  
30 human PSM gene is transferred to the rat tumor AT-6. AT-6 is a metastatic androgen independent tumor. The same chromosome transferred into non prostate derived tissues or tumors is not expressed and thus these cells could be used

-81-

as an animal model for these experiments. PSA, PSM positive Human LNCaP cells will be used for testing in nude mice.

References of the Second Series of Experiments

1. Coffey, D.S. Prostate Cancer - An overview of an  
5 increasing dilemma. Cancer Supplement, 71,3: 880-886,  
1993.
2. Chiarodo, A. National Cancer Institute roundtable on  
prostate cancer; future research directions. Cancer  
10 Res., 51: 2498-2505, 1991.
3. Israeli, R.S., Powell, C.T., Fair, W.R., and Heston,  
W.D.W. Molecular cloning of a complementary DNA  
encoding a prostate-specific membrane antigen. Cancer  
15 Res., 53: 227-230, 1993.
4. Horoszewicz, J.S., Kawinski, E., and Murphy, G.P.  
Monoclonal antibodies to a new antigenic marker in  
epithelial cells and serum of prostatic cancer  
20 patients. Anticancer Res., 7: 927-936, 1987.
5. Horoszewicz, J.S., Leong, S.S., Kawinski, E., Karr,  
J.P., Rosenthal, H., Chu, T.M., Mirand, E.A., and  
Murphy, G.P. LNCaP model of human prostatic  
25 carcinoma. Cancer Res., 43: 1809-1818, 1983.
6. Abdel-Nabi, H., Wright, G.L., Gulfo, J.V., Petrylak,  
D.P., Neal, C.E., Texter, J.E., Begun, F.P., Tyson,  
I., Heal, A., Mitchell, E., Purnell, G., and Harwood,  
30 S.J. Monoclonal antibodies and radioimmunoconjugates  
in the diagnosis and treatment of prostate cancer.  
Semin. Urol., 10: 45-54, 1992.
7. Stone, K.R., Mickey, D.D., Wunderli, H., Mickey, G.H.,  
35 and Paulson, D.F. Isolation of a human prostate

-83-

carcinoma cell line (DU-145). Int. J. Cancer, 21: 274-281, 1978.

- 5 8. Kaign, M.E., Narayan, K.S., Ohnuki, Y., and Lechner, J.F. Establishment and characterization of a human prostatic carcinoma cell line (PC-3). Invest. Urol., 17: 16-23, 1979.
- 10 9. Hsu, S.M., Raine, L., and Fanger, H. Review of present methods of immunohistochemical detection. Am. J. Clin. Path. 75: 734-738, 1981.
- 15 10. Harlow, E., and Lane, D. Antibodies: A Laboratory Manual. New York: Cold Spring Harbor Laboratory, p. 449, 1988.
- 20 11. Glisin, V., Crkvenjakov, R., and Byus, C. Ribonucleic acid isolated by cesium chloride centrifugation. Biochemistry, 13: 2633-2637, 1974.
- 25 12. Aviv, H., and Leder, P. Purification of biologically active globin messenger RNA by chromatography on oligo-thymidylic acid cellulose. Proc. Natl. Acad. Sci. USA, 69: 1408-1412, 1972.
- 30 13. Melton, D.A., Krieg, P.A., Rebagliati, M.R., Maniatis, T.A., Zinn, K., and Careen, M.R. Efficient in-vitro synthesis of biologically active RNA and RNA hybridization probes from plasmids containing a bacteriophage SP6 promoter. Nucl. Acids. Res. 12: 7035-7056, 1984.
- 35 14. Personal Communication from Cytogen Corporation, Princeton, N.J.

- 84 -

15. Axelrod, H.R., Gilman, S.C., D'Aleo, C.J., Petrylak, D., Reuter, V., Gulfo, J.V., Saad, A., Cordon-Cardo, C., and Scher, H.I. Preclinical results and human immunohistochemical studies with <sup>90</sup>Y-CYT-356; a new  
5 prostatic cancer therapeutic agent. AUA Proceedings, Abstract 596, 1992.
16. Lopes, A.D., Davis, W.L., Rosenstraus, M.J., Uveges, A.J., and Gilman, S.C. Immunohistochemical and  
10 pharmacokinetic characterization of the site-specific immunoconjugate CYT-356 derived from antiprostata monoclonal antibody 7E11-C5. Cancer Res., 50: 6423-6429, 1990.
- 15 17. Troyer, J.K., Qi, F., Beckett, M.L., Morningstar, M.M., and Wright, G.L. Molecular characterization of the 7E11-C5 prostate tumor-associated antigen. AUA Proceedings. Abstract 482, 1993.
- 20 18. Roemer, K., Friedmann, T. Concepts and strategies for human gene therapy. FEBS. 223:212-225.
19. Antonie, P. Springer, C.J., Bagshawe, F., Searle, F., Melton, R.G., Rogers, G.T., Burke, P.J., Sherwood,  
25 R.F. Disposition of the prodrug 4-bis(2chloroethyl) amino) benzoyl-1-glutamic acid and its active parent drug in mice. Br.J.Cancer 62:909-914, 1990.
20. Connor, J. Bannerji, R., Saito, S., Heston, W.D.W.,  
30 Fair, W.R., Gilboa, E. Regression of bladder tumors in mice treated with interleukin 2 gene-modified tumor cells. J.Exp.Med. 177:1127-1134, 1993. (appendix)
21. Vile R., Hart, I.R. In vitro and in vivo targeting of

-85-

gene expression to melanoma cells. Cancer Res. 53:962-967, 1993.

- 5 22. Warner, J.A., Heston, W.D.W. Future developments of nonhormonal systemic therapy for prostatic carcinoma. Urologic Clinics of North America 18:25-33, 1991.
- 10 23. Vile, R.G., Hart, I.R. Use of tissue specific expression of the herpes simplex virus thymidine kinase gene to inhibit growth of established murine melanomas following direct intratumoral injection of DNA. Cancer Res. 53:3860-3864, 1993.

-86-

Third Series of ExperimentsSensitive Detection of Prostatic Hematogenous  
Micrometastases Using PSA and PSM-Derived Primers in the  
Polymerase Chain Reaction

We have developed a PCR-based assay enabling sensitive detection of hematogenous micrometastases in patients with prostate cancer. We performed "nested PCR", amplifying mRNA sequences unique to prostate-specific antigen and to the prostate-specific membrane antigen, and have compared their respective results. Micrometastases were detected in 2/30 patients (6.7%) by PCR with PSA-derived primers, while PSM-derived primers detected tumor cells in 19/16 patients (63.3%). All 8 negative controls were negative with both PSA and PSM PCR. Assays were repeated to confirm results, and PCR products were verified by DNA sequencing and Southern analysis. Patients harboring circulating prostatic tumor cells as detected by PSM, and not by PSA-PCR included 4 patients previously treated with radical prostatectomy and with non-measurable serum PSA levels at the time of this assay. The significance of these findings with respect to future disease recurrence and progression will be investigated.

Improvement in the overall survival of patients with prostate cancer will depend upon earlier diagnosis. Localized disease, without evidence of extra-prostatic spread, is successfully treated with either radical prostatectomy or external beam radiation, with excellent long-term results (2,3). The major problem is that approximately two-thirds of men diagnosed with prostate cancer already have evidence of advanced extra-prostatic spread at the time of diagnosis, for which there is at present no cure (4). The use of clinical serum markers



-87-

such as prostate-specific antigen (PSA) and prostatic acid phosphatase (PAP) have enabled clinicians to detect prostatic carcinomas earlier and provide useful parameters to follow responses to therapy (5). Yet, despite the advent  
5 of sensitive serum PSA assays, radionuclide bone scans, CT scans and other imaging modalities, we are still unable to detect the presence of micrometastatic cells prior to their establishment of solid metastases. Previous work has been done utilizing the polymerase chain reaction to amplify mRNA  
10 sequences unique to breast, leukemia, and other malignant cells in the circulation and enable early detection of micrometastases (6,7). Recently, a PCR-based approach utilizing primers derived from the PSA DNA sequence was published (8). In this study 3/12 patients with advanced,  
15 stage D prostate cancer had detectable hematogenous micrometastases.

We have recently identified and cloned a 2.65 kb cDNA encoding the 100 kDa prostate-specific membrane antigen  
20 (PSM) recognized by the anti-prostate monoclonal antibody 7E11-C5.3 (9). PSM appears to be an integral membrane glycoprotein which is very highly expressed in prostatic tumors and metastases and is almost entirely prostate-specific (10). Many anaplastic tumors and bone metastases  
25 have variable and at times no detectable expression of PSA, whereas these lesions appear to consistently express high levels of PSM. Prostatic tumor cells that escape from the prostate gland and enter the circulation are likely to have the potential to form metastases and are possibly the more  
30 aggressive and possibly anaplastic cells, a population of cells that may not express high levels of PSA, but may retain high expression of PSM. We therefore chose to utilize DNA primers derived from the sequences of both PSA and PSM in a PCR assay to detect micrometastatic cells in  
35 the peripheral circulation. Despite the high level of

- 88 -

amplification and sensitivity of conventional RNA PCR, we have utilized a "nested" PCR approach in which we first amplify a target sequence, and subsequently use this PCR product as the template for another round of PCR amplification with a new set of primers totally contained within the sequence of the previous product. This approach has enabled us to increase our level of detection from one prostatic tumor cell per 10,000 cells to better than one cell per ten million cells.

10 --

### Experimental Details

#### Materials and Methods

##### 15 Cells and Reagents:

15 LNCaP and MCF-7 cells were obtained from the American Type Culture Collection (Rockville, MD.). Details regarding the establishment and characteristics of these cell lines have been previously published (11,12). Cells were grown in RPMI  
20 1640 media supplemented with L-glutamine, nonessential amino acids, obtained from the MSKCC Media Preparation Facility, and 5% fetal calf serum (Gibco-BRL, Gaithersburg, MD.) in a CO<sub>2</sub> incubator at 37C. All cell media was obtained from the MSKCC Media Preparation Facility. Routine chemical reagents  
25 were of the highest grade possible and were obtained from Sigma Chemical Company, St. Louis, MO.

##### Patient Blood Specimens

30 All blood specimens used in this study were from patients seen in the outpatient offices of urologists on staff at MSKCC. Two anti-coagulated (purple top) tubes per patient were obtained at the time of their regularly scheduled blood draws. Specimen procurement was conducted as per the approval of the MSKCC Institutional Review Board. Samples

- 89 -

were promptly brought to the laboratory for immediate processing. Serum PSA and PAP determinations were performed by standard techniques by the MSKCC Clinical Chemistry Laboratory. PSA determinations were performed using the  
5 Tandem PSA assay (Hybritech, San Diego, CA.). The eight blood specimens used as negative controls were from 2 males with normal serum PSA values and biopsy-proven BPH, one healthy female, 3 healthy males, one patient with bladder cancer, and one patient with acute promyelocytic leukemia.

10

#### Blood Sample Processing/RNA Extraction

4 ml of whole anticoagulated venous blood was mixed with 3 ml of ice cold phosphate buffered saline and then carefully layered atop 8 ml of Ficoll (Pharmacia, Uppsala, Sweden) in  
15 a 15-ml polystyrene tube. Tubes were centrifuged at 200 x g for 30 min. at 4C. Using a sterile pasteur pipette, the buffy coat layer (approx. 1 ml.) was carefully removed and rediluted up to 50 ml with ice cold phosphate buffered saline in a 50 ml polypropylene tube. This tube was then  
20 centrifuged at 2000 x g for 30 min at 4C. The supernatant was carefully decanted and the pellet was allowed to drip dry. One ml of RNazol B was then added to the pellet and total RNA was isolated as per manufacturers directions (Cinna/Biotechx, Houston, TX.). RNA concentrations and  
25 purity were determined by UV spectroscopy on a Beckman DU 640 spectrophotometer and by gelanalysis.

#### Determination of PCR Sensitivity

RNA was isolated from LNCaP cells and from mixtures of LNCaP  
30 and MCF-7 cells at fixed ratios (i.e. 1:100, 1:1000, etc.) using RNazol B. Nested PCR was then performed as described below with both PSA and PSM primers in order to determine the limit of detection for the assay. LNCaP:MCF-7 (1:100,000) cDNA was diluted with distilled water to obtain  
35 concentrations of 1:1,000,000 and 1:10,000,000. MCF-7 cells

-90-

were chosen because they have been previously tested and shown not to express PSM by PCR.

#### Polymerase Chain Reaction

5 The PSA outer primers used span portions of exons 4 and 5 to yield a 486 bp PCR product and enable differentiation between cDNA and possible contaminating genomic DNA amplification. The upstream primer sequence beginning at nucleotide 494 in PSA cDNA sequence is 5'-  
10 TACCCACTGCATCAGGAACA-3' (SEQ. ID. No. 39) and the downstream primer at nucleotide 960 is 5'-CCTTGAAGCACACCATTACA-3' (SEQ. ID. No. 40). The PSA inner upstream primer (beginning at nucleotide 559) 5'-ACACAGGCCAGGTATTTTCAG-3' (SEQ. ID. No. 41) and the downstream primer (at nucleotide 894) 5'-  
15 GTCCAGCGTCCAGCACACAG-3' (SEQ. ID. No. 42) yield a 355 bp PCR product. All primers were synthesized by the MSKCC Microchemistry Core Facility. 5g of total RNA was reverse-transcribed into cDNA in a total volume of 20l using Superscript reverse transcriptase (Gibco-BRL) according to  
20 the manufacturers recommendations. 1l of this cDNA served as the starting template for the outer primer PCR reaction. The 20l PCR mix included: 0.5U Taq polymerase (Promega Corp., Madison, WI.), Promega reaction buffer, 1.5mM MgCl<sub>2</sub>, 200M dNTPs, and 1.0M of each primer. This mix was then  
25 transferred to a Perkin Elmer 9600 DNA thermal cycler and incubated for 25 cycles. The PCR profile was as follows: 94C x 15 sec., 60C x 15 sec., and 72C for 45 sec. After 25 cycles, samples were placed on ice, and 1l of this reaction mix served as the template for another round of PCR using  
30 the inner primers. The first set of tubes were returned to the thermal cycler for 25 additional cycles. PSM-PCR required the selection of primer pairs that also spanned an intron in order to be certain that cDNA and not genomic DNA were being amplified. Since the genomic DNA sequence of PSM

-91-

has not yet been determined, this involved trying different primer pairs until a pair was found that produced the expected size PCR product when cDNA was amplified, but with no band produced from a genomic DNA template, indicating the presence of a large intron. The PSM outer primers yield a 946 bp product and the inner primers a 434 bp product. The PSM outer upstream primer used was 5'-ATGGGTGTTTGGTGGTATTGACC-3' (SEQ. ID. No. 43) (beginning at nucleotide 1401) and the downstream primer (at nucleotide 2348) was 5'-TGCTTGGAGCATAGATGACATGC-3' (SEQ. ID. No. 44). The PSM inner upstream primer (at nucleotide 1581) was 5'-ACTCCTTCAAGAGCGTGGCG-3' (SEQ. ID. No. 45) and the downstream primer (at nucleotide 2015) was 5'-AACACCATCCCTCCTCGAACC-3' (SEQ. ID. No. 46). cDNA used was the same as for the PSA assay. The 50l PCR mix included: 1U Taq Polymerase (Promega), 250M dNTPs, 10mM -mercaptoethanol, 2mM MgCl<sub>2</sub>, and 5l of a 10x buffer mix containing: 166mM NH<sub>4</sub>SO<sub>4</sub>, 670mM Tris pH 8.8, and 2 mg/ml of acetylated BSA. PCR was carried out in a Perkin Elmer 480 DNA thermal cycler with the following parameters: 94C x 4 minutes for 1 cycle, 94C x 30 sec., 58C x 1 minute, and 72C x 1 minute for 25 cycles, followed by 72C x 10 minutes. Samples were then iced and 2l of this reaction mix was used as the template for another 25 cycles with a new reaction mix containing the inner PSM primers. cDNA quality was verified by performing control reactions using primers derived from -actin yielding a 446 bp PCR product. The upstream primer used was 5'-AGGCCAACCGCGAGAAGATGA-3' (SEQ. ID. No. 47) (exon 3) and the downstream primer was 5'-ATGTCACACTGGGGAAGC-3' (SEQ. ID. No. 48) (exon 4). The entire PSA mix and 10l of each PSM reaction mix were run on 1.5-2% agarose gels, stained with ethidium bromide and photographed in an Eagle Eye Video Imaging System (Stratagene, Torrey Pines, CA.). Assays were repeated at least 3 times to verify results.

- 92 -

Cloning and Sequencing of PCR Products

PCR products were cloned into the pCR II plasmid vector using the TA cloning system (Invitrogen). These plasmids were transformed into competent E. coli cells using standard  
5 methods (13) and plasmid DNA was isolated using Magic Minipreps (Promega) and screened by restriction analysis. TA clones were then sequenced by the dideoxy method (14) using Sequenase (U.S. Biochemical). 3-4g of each plasmid was denatured with NaOH and ethanol precipitated. Labeling  
10 reactions were carried out according to the manufacturers recommendations using <sup>35</sup>S-dATP (NEN), and the reactions were terminated as discussed in the same protocol. Sequencing products were then analyzed on 6% polyacrilamide/7M urea gels run at 120 watts for 2 hours. Gels were fixed for 20  
15 minutes in 10% methanol/10% acetic acid, transferred to Whatman 3MM paper and dried down in a vacuum dryer for 2 hours at 80C. Gels were then autoradiographed at room temperature for 18 hours.

20 Southern Analysis

Ethidium-stained agarose gels of PCR products were soaked for 15 minutes in 0.2N HCl, followed by 30 minutes each in 0.5N NaOH/1.5M NaCl and 0.1M Tris pH 7.5/1.5M NaCl. Gels were then equilibrated for 10 minutes in 10x SSC (1.5M  
25 NaCl/0.15M Sodium Citrate. DNA was transferred onto Nytran nylon membranes (Schleicher and Schuell) by pressure blotting in 10x SSC with a Posi-blotter (Stratagene). DNA was cross-linked to the membrane using a UV Stratalinker (Stratagene). Blots were pre-hybridized at 65C for 2 hours  
30 and subsequently hybridized with denatured <sup>32</sup>P-labeled, random-primed cDNA probes (either PSM or PSA) (9,15). Blots were washed twice in 1x SSPE/0.5% SDS at 42C and twice in 0.1x SSPE/0.5% SDS at 50C for 20 minutes each. Membranes were air-dried and autoradiographed for 30 minutes to 1 hour

-93-

at -70C with Kodak X-Omat film.

### Experimental Results

Our technique of PCR amplification with nested primers  
5 improved our level of detection of prostatic cells from  
approximately one prostatic cell per 10,000 MCF-7 cells to  
better than one cell per million MCF-7 cells, using either  
PSA or PSM-derived primers (figures 26 and 27). This  
represents a substantial improvement in our ability to  
10 detect minimal disease. Characteristics of the 16 patients  
analyzed with respect to their clinical stage, treatment,  
serum PSA and PAP values, and results of our assay are shown  
in table 1. In total, PSA-PCR detected tumor cells in 2/30  
patients (6.7%), whereas PSM-PCR detected cells in 19/30  
15 patients (63.3%). There were no patients positive for tumor  
cells by PSA and not by PSM, while PSM provided 8 positive  
patients not detected by PSA. Patients 10 and 11 in table  
1, both with very advanced hormone-refractory disease were  
detected by both PSA and PSM. Both of these patients have  
20 died since the time these samples were obtained. Patients  
4, 7, and 12, all of whom were treated with radical  
prostatectomies for clinically localized disease, and all of  
whom have non-measurable serum PSA values 1-2 years  
postoperatively were positive for circulating prostatic  
25 tumor cells by PSM-PCR, but negative by PSA-PCR. A  
representative ethidium stained gel photograph for PSM-PCR  
is shown in figure 28. Samples run in lane A represent PCR  
products generated from the outer primers and samples in  
lanes labeled B are products of inner primer pairs. The  
30 corresponding PSM Southern blot autoradiograph is shown in  
figure 29. The sensitivity of the Southern blot analysis  
exceeded that of ethidium staining, as can be seen in  
several samples where the outer product is not visible on  
figure 28, but is detectable by Southern blotting as shown  
35 in figure 29. In addition, sample 3 on figures 28 and 29

-94-

(patient 6 in figure 30) appears to contain both outer and inner bands that are smaller than the corresponding bands in the other patients. DNA sequencing has confirmed that the nucleotide sequence of these bands matches that of PSM, with the exception of a small deletion. This may represent either an artifact of PCR, alternative splicing of PSM mRNA in this patient, or a PSM mutation. We have noted similar findings with other samples on several occasions (unpublished data). All samples sequenced and analyzed by Southern analysis have been confirmed as true positives for PSA and PSM.

#### Experimental Details

The ability to accurately stage patients with prostate cancer at the time of diagnosis is clearly of paramount importance in selecting appropriate therapy and in predicting long-term response to treatment, and potential cure. Pre-surgical staging presently consists of physical examination, serum PSA and PAP determinations, and numerous imaging modalities including transrectal ultrasonography, CT scanning, radionuclide bone scans, and even MRI scanning. No present modality, however, addresses the issue of hematogenous micrometastatic disease and the potential negative impact on prognosis that this may produce. Previous work has shown that only a fractional percentage of circulating tumor cells will inevitably go on to form a solid metastasis (16), however, the detection of and potential quantification of circulating tumor cell burden may prove valuable in more accurately staging disease. The long-term impact of hematogenous micrometastatic disease must be studied by comparing the clinical courses of patients found to have these cells in their circulation with patients of similar stage and treatment who test negatively.

The significantly higher level of detection of tumor cells



- 95 -

with PSM as compared to PSA is not surprising to us, since we have noted more consistent expression of PSM in prostate carcinomas of all stages and grades as compared to variable expression of PSA in more poorly differentiated and anaplastic prostate cancers. We were surprised to detect tumor cells in the three patients that had undergone radical prostatectomies with subsequent undetectable amounts of serum PSA. These patients would be considered to be surgical "cures" by standard criteria, yet they apparently continue to harbor prostatic tumor cells. It will be interesting to follow the clinical course of these patients as compared to others without PCR evidence of residual disease. We are presently analyzing larger numbers of patient samples in order to verify these findings and perhaps identify patients at risk for metastatic disease.

References

1. Boring, C.C., Squires, T.S., and Tong, T.: Cancer Statistics, 1993. CA Cancer J. Clin., 43:7-26, 1993.
- 5 2. Lepor, H., and Walsh, P.C.: Long-term results of radical prostatectomy in clinically localized prostate cancer: Experience at the Johns Hopkins Hospital. NCI Monogr., 7:117-122, 1988.
- 10 3. Bagshaw, M.A., Cox, R.S., and Ray, G.R.: Status of radiation treatment of prostate cancer at Stanford University. NCI Monogr., 7:47-60, 1988.
- 15 4. Thompson, I.M., Rounder, J.B., Teague, J.L., et al.: Impact of routine screening for adenocarcinoma of the prostate on stage distribution. J. Urol., 137:424-426, 1987.
- 20 5. Chiarodo, A.: A National Cancer Institute roundtable on prostate cancer; future research directions. Cancer Res., 51:2498-2505, 1991.
- 25 6. Wu, A., Ben-Ezra, J., and Colombero, A.: Detection of micrometastasis in breast cancer by the polymerase chain reaction. Lab. Invest., 62:109A, 1990.
- 30 7. Fey, M.F., Kulozik, A.E., and Hansen-Hagge, T.E.: The polymerase chain reaction: A new tool for the detection of minimal residual disease in hematological malignancies. Eur. J. Cancer, 27:89-94, 1991.
- 35 8. Moreno, J.G., Croce, C.M., Fischer, R., Monne, M., Vihko, P., Mulholland, S.G., and Gomella, L.G.: Detection of hematogenous micrometastasis in patients

- 97 -

with prostate cancer. Cancer Res., 52:6110-6112, 1992.

- 5 9. Israeli, R.S., Powell, C.T., Fair, W.R., and Heston, W.D.W.: Molecular cloning of a complementary DNA encoding a prostate-specific membrane antigen. Cancer Res., 53:227-230, 1993.
- 10 10. Israeli, R.S., Powell, C.T., Corr, J.G., Fair, W.R., and Heston, W.D.W.: Expression of the prostate-specific membrane antigen (PSM): Submitted to Cancer Research.
- 15 11. Horoszewicz, J.S., Leong, S.S., Kawinski, E., Karr, J.P., Rosenthal, H., Chu, T.M., Mirand, E.A., and Murphy, G.P.: LNCaP model of human prostatic carcinoma. Cancer Res., 43:1809-1818, 1983.
- 20 12. Soule, H.D., Vazquez, J., Long, A., Albert, S., and Brennan, M.: A human cell line from a pleural effusion derived from a breast carcinoma. J. Natl. Can. Inst., 51:1409-1416, 1973.
- 25 13. Hanahan, D.: Studies on transformation of *Escherichia coli* with plasmids. J. Mol. Biol., 166:557-580, 1983.
- 30 14. Sanger, F., Nicklen, S., and Coulson, A.R.: DNA sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. USA, 74:5463-5467, 1977.
- 35 15. Lundwall, A., and Lilja, H.: Molecular cloning of a human prostate specific antigen cDNA. FEBS Letters, 214:317, 1987.

-98-

16. Liotta, L.A., Kleinerma, J., and Saidel, G.M.: Quantitative relationships of intravascular tumor cells, tumor vessels, and pulmonary metastases following tumor implantation. Cancer Res., 34:997-1003, 1974.
- 5

-99-

Fourth Series of Experiments5      EXPRESSION OF THE PROSTATE SPECIFIC MEMBRANE ANTIGEN (PSM)  
DIMINISHES THE MITOGENIC STIMULATION OF AGGRESSIVE HUMAN  
PROSTATIC CARCINOMA CELLS BY TRANSFERRIN

An association between transferrin and human prostate cancer has been suggested by several investigators. It has been  
10 shown that the expressed prostatic secretions of patients with prostate cancer are enriched with respect to their content of transferrin and that prostate cancer cells are rich in transferrin receptors (J. Urol. 143, 381, 1990). Transferrin derived from bone marrow has been shown to  
15 selectively stimulate the growth of aggressive prostate cancer cells (PNAS 89, 6197, 1992). We have previously reported the cloning of the cDNA encoding the 100 kDa PSM antigen (Cancer Res. 53, 208, 1993). DNA sequence analysis has revealed that a portion of the coding region, from  
20 nucleotide 1250 to 1700 possesses a 54% homology to the human transferrin receptor. PC-3 cells do not express PSM mRNA or protein and exhibit increased cell growth in response to transferrin, whereas, LNCaP prostate cancer cells which highly express PSM have a very weak response to  
25 transferrin. To determine whether PSM expression by prostatic cancer cells impacts upon their mitogenic response to transferrin we stably transfected the full-length PSM cDNA into the PC-3 prostate cancer cells. Clones highly

-100-

expressing PSM mRNA were identified by Northern analysis and expression of PSM protein was verified by Western analysis using the anti-PSM monoclonal antibody 7E11-C5.3.

5 We plated  $2 \times 10^4$  PC-3 or PSM-transfected PC-3 cells per well in RPMI medium supplemented with 10% fetal bovine serum and at 24 hrs. added 1  $\mu$ g per ml. of holotransferrin to the cells. Cells were counted at 1 day to be highly mitogenic to the PC-3 cells. Cells were counted at 1 day to determine  
10 plating efficiency and at 5 days to determine the effect of the transferrin. Experiments were repeated to verify the results.

We found that the PC-3 cells experienced an average increase  
15 of 275% over controls, whereas the LNCaP cells were only stimulated 43%. Growth kinetics revealed that the PSM-transfected PC-3 cells grew 30% slower than native PC-3 cells. This data suggests that PSM expression in aggressive, metastatic human prostate cancer cells  
20 significantly abrogates their mitogenic response to transferrin.

The use of therapeutic vaccines consisting of cytokine-secreting tumor cell preparations for the treatment of  
25 established prostate cancer was investigated in the Dunning R3327-MatLyLu rat prostatic adenocarcinoma model. Only IL-2

-101-

secreting, irradiated tumor cell preparations were capable of curing animals from subcutaneously established tumors, and engendered immunological memory that protected the animals from another tumor challenge. Immunotherapy was  
5 less effective when tumors were induced orthotopically, but nevertheless led to improved outcome, significantly delaying, and occasionally preventing recurrence of tumors after resection of the cancerous prostate. Induction of a  
10 potent immune response in tumor bearing animals against the nonimmunogenic MatLyLu tumor supports the view that active immunotherapy of prostate cancer may have therapeutic benefits.

- 102 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Israeli, Ron S.  
Heston, Warren D.W.  
Fair, William R.
- (ii) TITLE OF INVENTION: PROSTATE-SPECIFIC MEMBRANE ANTIGEN
- (iii) NUMBER OF SEQUENCES: 38
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Cooper & Dunham
  - (B) STREET: 30 Rockefeller Plaza
  - (C) CITY: New York
  - (D) STATE: New York
  - (E) COUNTRY: United States of America
  - (F) ZIP: 10112
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: White, John P.
  - (B) REGISTRATION NUMBER: 28,678
  - (C) REFERENCE/DOCKET NUMBER: 1747/41426
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (212) 977-9550
  - (B) TELEFAX: (212) 664-0525
  - (C) TELEX: 422523 COOP UI

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 2653 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapiens
  - (F) TISSUE TYPE: Carcinoma



-103-

## (vii) IMMEDIATE SOURCE:

(B) CLONE: Prostate-Specific Membrane Antigen

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 262..2511

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCAAAAGGG GCCGGATTTC CTTCTCCTGG AGGCAGATGT TGCCTCTCTC TCTCGCTCGG	60
ATTGGTTTCAG TGCACTCTAG AAACACTGCT GTGGTGGAGA AACTGGACCC CAGGTCTGGA	120
GCGAATTCCA GCCTGCAGGG CTGATAAGCG AGGCATTAGT GAGATTGAGA GAGACTTTAC	180
CCCCGCCGTGG TGGTTGGAGG GCGCGCAGTA GAGCAGCAGC ACAGGCGCGG GTCCCCGGGAG	240
GCCGGCTCTG CTCGCGCCGA G ATG TGG AAT CTC CTT CAC GAA ACC GAC TCG	291
Met Trp Asn Leu Leu His Glu Thr Asp Ser	
1 5 10	
GCT GTG GCC ACC GCG CGC CGC CCG CGC TGG CTG TGC GCT GGG GCG CTG	339
Ala Val Ala Thr Ala Arg Arg Pro Arg Trp Leu Cys Ala Gly Ala Leu	
15 20 25	
GTG CTG GCG GGT GGC TTC TTT CTC CTC GGC TTC CTC TTC GGG TGG TTT	387
Val Leu Ala Gly Gly Phe Phe Leu Leu Gly Phe Leu Phe Gly Trp Phe	
30 35 40	
ATA AAA TCC TCC AAT GAA GCT ACT AAC ATT ACT CCA AAG CAT AAT ATG	435
Ile Lys Ser Ser Asn Glu Ala Thr Asn Ile Thr Pro Lys His Asn Met	
45 50 55	
AAA GCA TTT TTG GAT GAA TTG AAA GCT GAG AAC ATC AAG AAG TTC TTA	483
Lys Ala Phe Leu Asp Glu Leu Lys Ala Glu Asn Ile Lys Lys Phe Leu	
60 65 70	
TAT AAT TTT ACA CAG ATA CCA CAT TTA GCA GGA ACA GAA CAA AAC TTT	531
Tyr Asn Phe Thr Gln Ile Pro His Leu Ala Gly Thr Glu Gln Asn Phe	
75 80 85 90	
CAG CTT GCA AAG CAA ATT CAA TCC CAG TGG AAA GAA TTT GGC CTG GAT	579
Gln Leu Ala Lys Gln Ile Gln Ser Gln Trp Lys Glu Phe Gly Leu Asp	
95 100 105	
TCT GTT GAG CTA GCA CAT TAT GAT GTC CTG TTG TCC TAC CCA AAT AAG	627
Ser Val Glu Leu Ala His Tyr Asp Val Leu Leu Ser Tyr Pro Asn Lys	
110 115 120	
ACT CAT CCC AAC TAC ATC TCA ATA ATT AAT GAA GAT GGA AAT GAG ATT	675
Thr His Pro Asn Tyr Ile Ser Ile Ile Asn Glu Asp Gly Asn Glu Ile	
125 130 135	
TTC AAC ACA TCA TTA TTT GAA CCA CCT CCT CCA GGA TAT GAA AAT GTT	723
Phe Asn Thr Ser Leu Phe Glu Pro Pro Pro Pro Gly Tyr Glu Asn Val	
140 145 150	
TCG GAT ATT GTA CCA CCT TTC AGT GCT TTC TCT CCT CAA GGA ATG CCA	771
Ser Asp Ile Val Pro Pro Phe Ser Ala Phe Ser Pro Gln Gly Met Pro	
155 160 165 170	

- 104 -

GAG	GGC	GAT	CTA	GTG	TAT	GTT	AAC	TAT	GCA	CGA	ACT	GAA	GAC	TTC	TTT	819
Glu	Gly	Asp	Leu	Val	Tyr	Val	Asn	Tyr	Ala	Arg	Thr	Glu	Asp	Phe	Phe	
			175						180					185		
AAA	TTG	GAA	CGG	GAC	ATG	AAA	ATC	AAT	TGC	TCT	GGG	AAA	ATT	GTA	ATT	867
Lys	Leu	Glu	Arg	Asp	Met	Lys	Ile	Asn	Cys	Ser	Gly	Lys	Ile	Val	Ile	
			190					195					200			
GCC	AGA	TAT	GGG	AAA	GTT	TTC	AGA	GGA	AAT	AAG	GTT	AAA	AAT	GCC	CAG	915
Ala	Arg	Tyr	Gly	Lys	Val	Phe	Arg	Gly	Asn	Lys	Val	Lys	Asn	Ala	Gln	
		205					210					215				
CTG	GCA	GGG	GCC	AAA	GGA	GTC	ATT	CTC	TAC	TCC	GAC	CCT	GCT	GAC	TAC	963
Leu	Ala	Gly	Ala	Lys	Gly	Val	Ile	Leu	Tyr	Ser	Asp	Pro	Ala	Asp	Tyr	
	220					225					230					
TTT	GCT	CCT	GGG	GTG	AAG	TCC	TAT	CCA	GAT	GGT	TGG	AAT	CTT	CCT	GGA	1011
Phe	Ala	Pro	Gly	Val	Lys	Ser	Tyr	Pro	Asp	Gly	Trp	Asn	Leu	Pro	Gly	
235					240					245					250	
GGT	GGT	GTC	CAG	CGT	GGA	AAT	ATC	CTA	AAT	CTG	AAT	GGT	GCA	GGA	GAC	1059
Gly	Gly	Val	Gln	Arg	Gly	Asn	Ile	Leu	Asn	Leu	Asn	Gly	Ala	Gly	Asp	
				255					260					265		
CCT	CTC	ACA	CCA	GGT	TAC	CCA	GCA	AAT	GAA	TAT	GCT	TAT	AGG	CGT	GGA	1107
Pro	Leu	Thr	Pro	Gly	Tyr	Pro	Ala	Asn	Glu	Tyr	Ala	Tyr	Arg	Arg	Gly	
			270					275					280			
ATT	GCA	GAG	GCT	GTT	GGT	CTT	CCA	AGT	ATT	CCT	GTT	CAT	CCA	ATT	GGA	1155
Ile	Ala	Glu	Ala	Val	Gly	Leu	Pro	Ser	Ile	Pro	Val	His	Pro	Ile	Gly	
		285					290					295				
TAC	TAT	GAT	GCA	CAG	AAG	CTC	CTA	GAA	AAA	ATG	GGT	GGC	TCA	GCA	CCA	1203
Tyr	Tyr	Asp	Ala	Gln	Lys	Leu	Leu	Glu	Lys	Met	Gly	Gly	Ser	Ala	Pro	
	300					305					310					
CCA	GAT	AGC	AGC	TGG	AGA	GGA	AGT	CTC	AAA	GTG	CCC	TAC	AAT	GTT	GGA	1251
Pro	Asp	Ser	Ser	Trp	Arg	Gly	Ser	Leu	Lys	Val	Pro	Tyr	Asn	Val	Gly	
315					320					325					330	
CCT	GGC	TTT	ACT	GGA	AAC	TTT	TCT	ACA	CAA	AAA	GTC	AAG	ATG	CAC	ATC	1299
Pro	Gly	Phe	Thr	Gly	Asn	Phe	Ser	Thr	Gln	Lys	Val	Lys	Met	His	Ile	
				335					340					345		
CAC	TCT	ACC	AAT	GAA	GTG	ACA	AGA	ATT	TAC	AAT	GTG	ATA	GGT	ACT	CTC	1347
His	Ser	Thr	Asn	Glu	Val	Thr	Arg	Ile	Tyr	Asn	Val	Ile	Gly	Thr	Leu	
			350					355					360			
AGA	GGA	GCA	GTG	GAA	CCA	GAC	AGA	TAT	GTC	ATT	CTG	GGA	GGT	CAC	CGG	1395
Arg	Gly	Ala	Val	Glu	Pro	Asp	Arg	Tyr	Val	Ile	Leu	Gly	Gly	His	Arg	
			365				370					375				
GAC	TCA	TGG	GTG	TTT	GGT	GGT	ATT	GAC	CCT	CAG	AGT	GGA	GCA	GCT	GTT	1443
Asp	Ser	Trp	Val	Phe	Gly	Gly	Ile	Asp	Pro	Gln	Ser	Gly	Ala	Ala	Val	
	380					385					390					
GTT	CAT	GAA	ATT	GTG	AGG	AGC	TTT	GGA	ACA	CTG	AAA	AAG	GAA	GGG	TGG	1491
Val	His	Glu	Ile	Val	Arg	Ser	Phe	Gly	Thr	Leu	Lys	Lys	Glu	Gly	Trp	
395					400					405					410	
AGA	CCT	AGA	AGA	ACA	ATT	TTG	TTT	GCA	AGC	TGG	GAT	GCA	GAA	GAA	TTT	1539
Arg	Pro	Arg	Arg	Thr	Ile	Leu	Phe	Ala	Ser	Trp	Asp	Ala	Glu	Glu	Phe	

- 105 -

415										420					425					
GGT	CTT	CTT	GGT	TCT	ACT	GAG	TGG	GCA	GAG	GAG	AAT	TCA	AGA	CTC	CTT	1587				
Gly	Leu	Leu	Gly	Ser	Thr	Glu	Trp	Ala	Glu	Glu	Asn	Ser	Arg	Leu	Leu					
			430						435				440							
CAA	GAG	CGT	GGC	GTG	GCT	TAT	ATT	AAT	GCT	GAC	TCA	TCT	ATA	GAA	GGA	1635				
Gln	Glu	Arg	Gly	Val	Ala	Tyr	Ile	Asn	Ala	Asp	Ser	Ser	Ile	Glu	Gly					
		445					450					455								
AAC	TAC	ACT	CTG	AGA	GTT	GAT	TGT	ACA	CCG	CTG	ATG	TAC	AGC	TTG	GTA	1683				
Asn	Tyr	Thr	Leu	Arg	Val	Asp	Cys	Thr	Pro	Leu	Met	Tyr	Ser	Leu	Val					
	460					465					470									
CAC	AAC	CTA	ACA	AAA	GAG	CTG	AAA	AGC	CCT	GAT	GAA	GGC	TTT	GAA	GGC	1731				
His	Asn	Leu	Thr	Lys	Glu	Leu	Lys	Ser	Pro	Asp	Glu	Gly	Phe	Glu	Gly					
					480					485					490					
AAA	TCT	CTT	TAT	GAA	AGT	TGG	ACT	AAA	AAA	AGT	CCT	TCC	CCA	GAG	TTC	1779				
Lys	Ser	Leu	Tyr	Glu	Ser	Trp	Thr	Lys	Lys	Ser	Pro	Ser	Pro	Glu	Phe					
				495					500					505						
AGT	GGC	ATG	CCC	AGG	ATA	AGC	AAA	TTG	GGA	TCT	GGA	AAT	GAT	TTT	GAG	1827				
Ser	Gly	Met	Pro	Arg	Ile	Ser	Lys	Leu	Gly	Ser	Gly	Asn	Asp	Phe	Glu					
			510					515					520							
GTG	TTC	TTC	CAA	CGA	CTT	GGA	ATT	GCT	TCA	GGC	AGA	GCA	CGG	TAT	ACT	1875				
Val	Phe	Phe	Gln	Arg	Leu	Gly	Ile	Ala	Ser	Gly	Arg	Ala	Arg	Tyr	Thr					
		525					530					535								
AAA	AAT	TGG	GAA	ACA	AAC	AAA	TTC	AGC	GGC	TAT	CCA	CTG	TAT	CAC	AGT	1923				
Lys	Asn	Trp	Glu	Thr	Asn	Lys	Phe	Ser	Gly	Tyr	Pro	Leu	Tyr	His	Ser					
		540				545					550									
GTC	TAT	GAA	ACA	TAT	GAG	TTG	GTG	GAA	AAG	TTT	TAT	GAT	CCA	ATG	TTT	1971				
Val	Tyr	Glu	Thr	Tyr	Glu	Leu	Val	Glu	Lys	Phe	Tyr	Asp	Pro	Met	Phe					
	555				560					565					570					
AAA	TAT	CAC	CTC	ACT	GTG	GCC	CAG	GTT	CGA	GGA	GGG	ATG	GTG	TTT	GAG	2019				
Lys	Tyr	His	Leu	Thr	Val	Ala	Gln	Val	Arg	Gly	Gly	Met	Val	Phe	Glu					
				575					580					585						
CTA	GCC	AAT	TCC	ATA	GTG	CTC	CCT	TTT	GAT	TGT	CGA	GAT	TAT	GCT	GTA	2067				
Leu	Ala	Asn	Ser	Ile	Val	Leu	Pro	Phe	Asp	Cys	Arg	Asp	Tyr	Ala	Val					
			590					595					600							
GTT	TTA	AGA	AAG	TAT	GCT	GAC	AAA	ATC	TAC	AGT	ATT	TCT	ATG	AAA	CAT	2115				
Val	Leu	Arg	Lys	Tyr	Ala	Asp	Lys	Ile	Tyr	Ser	Ile	Ser	Met	Lys	His					
		605					610					615								
CCA	CAG	GAA	ATG	AAG	ACA	TAC	AGT	GTA	TCA	TTT	GAT	TCA	CTT	TTT	TCT	2163				
Pro	Gln	Glu	Met	Lys	Thr	Tyr	Ser	Val	Ser	Phe	Asp	Ser	Leu	Phe	Ser					
		620				625					630									
GCA	GTA	AAG	AAT	TTT	ACA	GAA	ATT	GCT	TCC	AAG	TTC	AGT	GAG	AGA	CTC	2211				
Ala	Val	Lys	Asn	Phe	Thr	Glu	Ile	Ala	Ser	Lys	Phe	Ser	Glu	Arg	Leu					
	635				640					645					650					
CAG	GAC	TTT	GAC	AAA	AGC	AAC	CCA	ATA	GTA	TTA	AGA	ATG	ATG	AAT	GAT	2259				
Gln	Asp	Phe	Asp	Lys	Ser	Asn	Pro	Ile	Val	Leu	Arg	Met	Met	Asn	Asp					
				655					660					665						

-106-

CAA CTC ATG TTT CTG GAA AGA GCA TTT ATT GAT CCA TTA GGG TTA CCA Gln Leu Met Phe Leu Glu Arg Ala Phe Ile Asp Pro Leu Gly Leu Pro 670 675 680	2307
GAC AGG CCT TTT TAT AGG CAT GTC ATC TAT GCT CCA AGC AGC CAC AAC Asp Arg Pro Phe Tyr Arg His Val Ile Tyr Ala Pro Ser Ser His Asn 685 690 695	2355
AAG TAT GCA GGG GAG TCA TTC CCA GGA ATT TAT GAT GCT CTG TTT GAT Lys Tyr Ala Gly Glu Ser Phe Pro Gly Ile Tyr Asp Ala Leu Phe Asp 700 705 710	2403
ATT GAA AGC AAA GTG GAC CCT TCC AAG GCC TGG GGA GAA GTG AAG AGA Ile Glu Ser Lys Val Asp Pro Ser Lys Ala Trp Gly Glu Val Lys Arg 715 720 725 730	2451
CAG ATT TAT GTT GCA GCC TTC ACA GTG CAG GCA GCT GCA GAG ACT TTG Gln Ile Tyr Val Ala Ala Phe Thr Val Gln Ala Ala Ala Glu Thr Leu 735 740 745	2499
AGT GAA GTA GCC TAAGAGGATT CTTTAGAGAA TCCGTATTGA ATTTGTGTGG Ser Glu Val Ala 750	2551
TATGTCACCTC AGAAAGAATC GTAATGGGTA TATTGATAAA TTTTAAAATT GGTATATTTG	2611
AAATAAAGTT GAATATTATA TATAAAAAAA AAAAAAAAAA AA	2653

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 750 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Trp Asn Leu Leu His Glu Thr Asp Ser Ala Val Ala Thr Ala Arg 1 5 10 15
Arg Pro Arg Trp Leu Cys Ala Gly Ala Leu Val Leu Ala Gly Gly Phe 20 25 30
Phe Leu Leu Gly Phe Leu Phe Gly Trp Phe Ile Lys Ser Ser Asn Glu 35 40 45
Ala Thr Asn Ile Thr Pro Lys His Asn Met Lys Ala Phe Leu Asp Glu 50 55 60
Leu Lys Ala Glu Asn Ile Lys Lys Phe Leu Tyr Asn Phe Thr Gln Ile 65 70 75 80
Pro His Leu Ala Gly Thr Glu Gln Asn Phe Gln Leu Ala Lys Gln Ile 85 90 95
Gln Ser Gln Trp Lys Glu Phe Gly Leu Asp Ser Val Glu Leu Ala His 100 105 110
Tyr Asp Val Leu Leu Ser Tyr Pro Asn Lys Thr His Pro Asn Tyr Ile

-107-

115					120					125					
Ser	Ile	Ile	Asn	Glu	Asp	Gly	Asn	Glu	Ile	Phe	Asn	Thr	Ser	Leu	Phe
130						135					140				
Glu	Pro	Pro	Pro	Pro	Gly	Tyr	Glu	Asn	Val	Ser	Asp	Ile	Val	Pro	Pro
145					150					155					160
Phe	Ser	Ala	Phe	Ser	Pro	Gln	Gly	Met	Pro	Glu	Gly	Asp	Leu	Val	Tyr
				165					170					175	
Val	Asn	Tyr	Ala	Arg	Thr	Glu	Asp	Phe	Phe	Lys	Leu	Glu	Arg	Asp	Met
			180					185					190		
Lys	Ile	Asn	Cys	Ser	Gly	Lys	Ile	Val	Ile	Ala	Arg	Tyr	Gly	Lys	Val
	195						200					205			
Phe	Arg	Gly	Asn	Lys	Val	Lys	Asn	Ala	Gln	Leu	Ala	Gly	Ala	Lys	Gly
210						215					220				
Val	Ile	Leu	Tyr	Ser	Asp	Pro	Ala	Asp	Tyr	Phe	Ala	Pro	Gly	Val	Lys
225					230					235					240
Ser	Tyr	Pro	Asp	Gly	Trp	Asn	Leu	Pro	Gly	Gly	Gly	Val	Gln	Arg	Gly
				245					250					255	
Asn	Ile	Leu	Asn	Leu	Asn	Gly	Ala	Gly	Asp	Pro	Leu	Thr	Pro	Gly	Tyr
			260					265					270		
Pro	Ala	Asn	Glu	Tyr	Ala	Tyr	Arg	Arg	Gly	Ile	Ala	Glu	Ala	Val	Gly
		275					280					285			
Leu	Pro	Ser	Ile	Pro	Val	His	Pro	Ile	Gly	Tyr	Tyr	Asp	Ala	Gln	Lys
	290					295					300				
Leu	Leu	Glu	Lys	Met	Gly	Gly	Ser	Ala	Pro	Pro	Asp	Ser	Ser	Trp	Arg
305					310					315					320
Gly	Ser	Leu	Lys	Val	Pro	Tyr	Asn	Val	Gly	Pro	Gly	Phe	Thr	Gly	Asn
				325					330					335	
Phe	Ser	Thr	Gln	Lys	Val	Lys	Met	His	Ile	His	Ser	Thr	Asn	Glu	Val
			340					345					350		
Thr	Arg	Ile	Tyr	Asn	Val	Ile	Gly	Thr	Leu	Arg	Gly	Ala	Val	Glu	Pro
		355					360					365			
Asp	Arg	Tyr	Val	Ile	Leu	Gly	Gly	His	Arg	Asp	Ser	Trp	Val	Phe	Gly
	370					375					380				
Gly	Ile	Asp	Pro	Gln	Ser	Gly	Ala	Ala	Val	Val	His	Glu	Ile	Val	Arg
385					390					395					400
Ser	Phe	Gly	Thr	Leu	Lys	Lys	Glu	Gly	Trp	Arg	Pro	Arg	Arg	Thr	Ile
				405					410					415	
Leu	Phe	Ala	Ser	Trp	Asp	Ala	Glu	Glu	Phe	Gly	Leu	Leu	Gly	Ser	Thr
			420				425						430		
Glu	Trp	Ala	Glu	Glu	Asn	Ser	Arg	Leu	Leu	Gln	Glu	Arg	Gly	Val	Ala
		435					440					445			

-108-

Tyr Ile Asn Ala Asp Ser Ser Ile Glu Gly Asn Tyr Thr Leu Arg Val  
 450 455 460  
 Asp Cys Thr Pro Leu Met Tyr Ser Leu Val His Asn Leu Thr Lys Glu  
 465 470 475 480  
 Leu Lys Ser Pro Asp Glu Gly Phe Glu Gly Lys Ser Leu Tyr Glu Ser  
 485 490 495  
 Trp Thr Lys Lys Ser Pro Ser Pro Glu Phe Ser Gly Met Pro Arg Ile  
 500 505 510  
 Ser Lys Leu Gly Ser Gly Asn Asp Phe Glu Val Phe Phe Gln Arg Leu  
 515 520 525  
 Gly Ile Ala Ser Gly Arg Ala Arg Tyr Thr Lys Asn Trp Glu Thr Asn  
 530 535 540  
 Lys Phe Ser Gly Tyr Pro Leu Tyr His Ser Val Tyr Glu Thr Tyr Glu  
 545 550 555 560  
 Leu Val Glu Lys Phe Tyr Asp Pro Met Phe Lys Tyr His Leu Thr Val  
 565 570 575  
 Ala Gln Val Arg Gly Gly Met Val Phe Glu Leu Ala Asn Ser Ile Val  
 580 585 590  
 Leu Pro Phe Asp Cys Arg Asp Tyr Ala Val Val Leu Arg Lys Tyr Ala  
 595 600 605  
 Asp Lys Ile Tyr Ser Ile Ser Met Lys His Pro Gln Glu Met Lys Thr  
 610 615 620  
 Tyr Ser Val Ser Phe Asp Ser Leu Phe Ser Ala Val Lys Asn Phe Thr  
 625 630 635 640  
 Glu Ile Ala Ser Lys Phe Ser Glu Arg Leu Gln Asp Phe Asp Lys Ser  
 645 650 655  
 Asn Pro Ile Val Leu Arg Met Met Asn Asp Gln Leu Met Phe Leu Glu  
 660 665 670  
 Arg Ala Phe Ile Asp Pro Leu Gly Leu Pro Asp Arg Pro Phe Tyr Arg  
 675 680 685  
 His Val Ile Tyr Ala Pro Ser Ser His Asn Lys Tyr Ala Gly Glu Ser  
 690 695 700  
 Phe Pro Gly Ile Tyr Asp Ala Leu Phe Asp Ile Glu Ser Lys Val Asp  
 705 710 715 720  
 Pro Ser Lys Ala Trp Gly Glu Val Lys Arg Gln Ile Tyr Val Ala Ala  
 725 730 735  
 Phe Thr Val Gln Ala Ala Ala Glu Thr Leu Ser Glu Val Ala  
 740 745 750

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 8 amino acids  
 (B) TYPE: amino acid

-109-

- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapiens
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:  
Ser Leu Tyr Glu Ser Xaa Thr Lys  
1 5
- (2) INFORMATION FOR SEQ ID NO:4:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 15 amino acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: peptide
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (A) ORGANISM: Homo sapiens
    - (F) TISSUE TYPE: Carcinoma
  - (vii) IMMEDIATE SOURCE:
    - (B) CLONE: Prostate Specific Membrane Antigen
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:  
Ser Tyr Pro Asp Gly Xaa Asn Leu Pro Gly Gly Gly Val Gln Arg  
1 5 10 15
- (2) INFORMATION FOR SEQ ID NO:5:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 7 amino acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: peptide
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO

-110-

(vi) ORIGINAL SOURCE:  
    (A) ORGANISM: Homo sapien  
    (F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:  
    (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Phe Tyr Asp Pro Met Phe Lys  
1                            5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 9 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: double  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
    (A) ORGANISM: Homo sapien  
    (F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:  
    (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ile Tyr Asn Val Ile Gly Thr Leu Lys  
1                            5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 22 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: double  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
    (A) ORGANISM: Homo sapien  
    (F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:  
    (B) CLONE: Prostate Specific Membrane Antigen



-111-

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Phe Leu Tyr Xaa Xaa Thr Gln Ile Pro His Leu Ala Gly Thr Glu Gln  
1                      5                      10                      15  
Asn Phe Gln Leu Ala Lys  
                    20

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapien
- (F) TISSUE TYPE: Carcinoma

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Gly Val Ile Leu Tyr Ser Asp Pro Ala Asp Tyr Phe Ala Pro Asp Val  
1                      5                      10                      15  
Lys

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapien
- (F) TISSUE TYPE: Carcinoma

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

-112-

Pro Val Ile Leu Tyr Ser Asp Pro Ala Asp Tyr Phe Ala Pro Gly Val  
 1 5 10 15

Lys

## (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 15 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Ala Phe Ile Asp Pro Leu Gly Leu Pro Asp Arg Pro Phe Tyr Arg  
 1 5 10 15

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Tyr Ala Gly Glu Ser Phe Pro Gly Ile Tyr Asp Ala Leu Phe Asp Ile  
 1 5 10 15

Glu Ser Lys

-113-

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Thr Ile Leu Phe Ala Ser Trp Asp Ala Glu Glu Phe Gly Xaa Xaa Glu  
1                    5                    10                    15  
Ser Thr Glu Glu Ala Glu  
                    20

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TTYTAYGAYC CNATGTT

17

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double

- 114 -

- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AACATNGGRT CRTARAA

17

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ATHTAYAAYG TNATHGG

17

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien

-115-

(F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:

(B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CCDATNACRT TRTADAT

17

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapien

(F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:

(B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CCNGCNGAYT AYTTCG

17

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapien

(F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:

(B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GCRAARTART CNGCNGG

17

-116-

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ACNGARCARA AYTTCARCT

20

## (2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

AGYTGRAART TYTGTCNGT

20

## (2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA

-117-

- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GARCARAAYT TYCARCT

17

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

AGYTGRAART TYTGYTC

17

## (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

-118-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TGGGAYGCNG ARGARTTYGG

20

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapien
- (F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:

- (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCRAAYTCYT CNGCRTCCCA

20

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapien
- (F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:

- (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TGGGAYGCNG ARGARTT

17

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid



- 119 -

(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapien
  - (F) TISSUE TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

AAATCYTCNG CRTCCCA

17

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 780 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

TACACTTATC CCATTTCGGAC ATGCCCCACCT TGGAAGTGA GACCCTTACA CCCCAGGCTT	60
CCCTTCGTTT AACCACACCC ANNNGTTTCC ACCAGTTGAA TCTTCAGGAC TACCCACAT	120
TGCTGTTT CAG ACCATCTCTA GCAGTGCAGC AGCCAGGCTG TTCAGCAAAA TGGATGGAGA	180
CACATGCTCT GANAGNNGTT GGAAAGGTGC GATCCANNNT TCCTGTAAGG TNNGACNNAA	240
CAAAGCAGGA GANNNGCCA GANTAATGGT GAAACTAGAT GTGAACAATT CCATGAAAGA	300
CAGGAAGATT CTGAACATCT TCGGTGCTAT CCAGGGATTT GAAGAACCTG ATCGGTATGT	360
TGTGATTGGA GCCCAGAGAG ACTCCTGGGG CCCAGGAGTG GCTAAAGCTG GCACTGGAAC	420
TGCTATATTG TTGGAAGTTG CCCGTGTGAT CTCAGACATA GTGAAAAACG AGGGCTACAA	480
ACCGAGGCGA AGCATCATCT TTGCTAGCTG GAGTGCAGGA GACTACGGAG CTGTGGGTGC	540
TACTGAATGG CTGGAGGGGT ACTCTGCCAT GCTGCATGCC AAAGCTTTCA CTTACATCAN	600
NGCTTGGATG CTCCAGTCCT GGGAGCAAGC CATGTCAAGA TTTCTGCCAG CCCCTTGCTG	660
TATATGCTGC TGGGGAGTAT TATGAAGGGG GTGAAGAATC CAGCAGCAGT CTCAGAGAGC	720

-120-

NNNNCTCTAT AACAGACTTG GCCCAGACTG GGTAAAAGCA GTTGTTCCTC TTGGCCTGGA 780

## (2) INFORMATION FOR SEQ ID NO:28:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 660 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

TGCAGAAAAG CTATTCAAAA ACATGGAAGG AAAGTGTCTT CCTAGTTGGA ATATAGATTC	60
CTCATGTAAG CTGGAACCTT CACAGAATCA AAATGTGAAG CTCAGTGTGA ACAATGTACT	120
GAAAGAAACA AGAATACTTA ACATCTTTGG CGTTATTAAA GGCTATGAGG AACCAGACCG	180
CTACATTGTA GTAGGAGCCC AGAGAGACGC TTGGGGCCCT GGTNGTTGCG AAGTCCAGTG	240
TGGGAACAGG TCTTNTCTGT GAAAGTTGCC CAAGTATTCT CAGATATGAT TTCAAAAGAT	300
GGATTTAGAC CCAGCAGGAG TATTATCTTT GCCAGCTGGA CTGCAGGAGA CTATGGAGCT	360
GTTGGTCCGA CTGAGTGGCT GGAGGGGTAC CTTTCATCTT TGCATCTAAA GNNNGCTTTC	420
ACTTACATTA ATNCTGGATA AAGTCGTCCT GGGTACTAGC AACTTCAAGG TTTCTGCCAG	480
CCCCCTATTA TATACACTTA TGGGGAAGAT AATGCAGGAN NCGTAAAGCA TCCGANNNNN	540
NNNTTGATGG AAAATATCTA TATCGAAACA GTAATTGGAT TAGCAAAATT GAGGAACCTT	600
CCTTGGACAA TGCTGCATTC CCTTTTCTTG CATATTCAGG AATCCCAGCA GTTTCTTTCT	660

## (2) INFORMATION FOR SEQ ID NO:29:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 540 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

- 121 -

TATGGAAGGA GACTGTCCCT CTGACTGGAA AACAGACTCT ACATGTAGGA TGGTAACCTC	60
AGAAAGCAAG AATGTGAAGC TCACTGTGAG CAATGTGCTG AAAGAGATAA AAATTCTTAA	120
CATCTTTGGA GTTATTAAAG GCTTTGTAGA ACCAGATCAC TATGTTGTAG TTGGGGCCCA	180
GAGAGATGCA TGGGGCCCTG GAGCTGCAAA ATCNCGGTGT AGGCACAGCT CTCCTATTGA	240
AACTTGCCCA GATGTTCTCA GATATGGTCT TAAAAGATGG GTTTCAGCCC AGCAGAAGCA	300
TTATCTTTGC CAGTTGGAGT GCTGGAGACT TTGGATCGGT TGGTGCCACT GAATGGCTAG	360
AGGGATACCT TTCGTCNCCT GCATTTAAAG GCTTTCACCT ATATTAATCT GGATAAAGCG	420
GTTCTTGGA CCAGCAACTT CAAGGTTTCT GCCAGCCAC TGTGTATAC GCTTATTGAG	480
AAAACAATGC AAAATGTGAA GCATCCGGT ACTGGGCAAT TTCTATATCA GGACAGCAAC	540

## (2) INFORMATION FOR SEQ ID NO:30:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapien
- (F) TISSUE TYPE: Carcinoma

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: Prostate Specific Membrane Antigen

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

ACGGAGCAAA ACTTTCAGCT TGCAAAG

27

## (2) INFORMATION FOR SEQ ID NO:31:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapien
- (F) TISSUE TYPE: Carcinoma

-122-

(vii) IMMEDIATE SOURCE:  
(B) CLONE: Prostate Membrane Specific Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Thr Glu Gln Asn Phe Gln Leu Ala Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 36 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapien  
(F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:  
(B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CTCTTCGGCA TCCCAGCTTG CAAACAAAAT TGTTCCT

36

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 36 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapien  
(F) TISSUE TYPE: Carcinoma

(vii) IMMEDIATE SOURCE:  
(B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

AGAACAATTT TGTTCGCAAG CTGGGATGCC AAGGAG

36

(2) INFORMATION FOR SEQ ID NO:34:

-123-

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 12 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapien
  - (G) CELL TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Arg Thr Ile Leu Phe Ala Ser Trp Asp Ala Glu Glu  
1                    5                    10

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 6 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapien
  - (G) CELL TYPE: Carcinoma
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Asp Glu Leu Lys Ala Glu  
1                    5

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 6 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide

-124-

- (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (A) ORGANISM: Homo Sapien
    - (G) CELL TYPE: Carcinoma
  - (vii) IMMEDIATE SOURCE:
    - (B) CLONE: Prostate Specific Membrane Antigen
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:  
Asn Glu Asp Gly Asn Glu  
1 5
- (2) INFORMATION FOR SEQ ID NO:37:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 6 amino acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: peptide
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (A) ORGANISM: Homo Sapien
    - (G) CELL TYPE: Carcinoma
  - (vii) IMMEDIATE SOURCE:
    - (B) CLONE: Prostate Specific Membrane Antigen
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:  
Lys Ser Pro Asp Glu Gly  
1 5
- (2) INFORMATION FOR SEQ ID NO:38:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 17 amino acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: peptide
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (A) ORGANISM: Homo Sapien
    - (G) CELL TYPE: Carcinoma

-125-

(vii) IMMEDIATE SOURCE:

(B) CLONE: Prostate Specific Membrane Antigen

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Ala Gly Ala Leu Val Leu Ala Gly Gly Phe Phe Leu Leu Gly Phe Leu  
1                      5                      10                      15

Phe

-126-

What is claimed is:

1. An isolated mammalian nucleic acid molecule encoding a mammalian prostate-specific membrane antigen.  
5
2. An isolated mammalian DNA molecule of claim 1.
3. An isolated mammalian cDNA molecule of claim 2.  
10
4. An isolated mammalian RNA molecule of claim 1.
5. An isolated mammalian nucleic acid molecule of claim 3, wherein the nucleic acid molecule is derived from humans.  
15
6. A nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence of the nucleic acid molecule of claim 1.  
20
7. A DNA molecule of claim 6.
8. An RNA molecule of claim 6.
9. A nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence of a nucleic acid molecule which is complementary to the nucleic acid molecule of claim 1.  
25
10. A DNA molecule of claim 9.  
30
11. An RNA molecule of claim 9.
12. A method of detecting expression of a mammalian prostate-specific membrane antigen in a cell which  
35



-127-

- 5 comprises obtaining total mRNA from the cell, contacting the mRNA so obtained with a labelled nucleic acid molecule of claim 6 under hybridizing conditions, determining the presence of mRNA hybridized to the molecule, and thereby detecting the expression of the mammalian prostate-specific membrane antigen in the cell.
- 10 13. A method of detecting expression of a mammalian prostate-specific membrane antigen in tissue sections which comprises contacting the tissue sections with a labelled nucleic acid molecule of claim 6 under hybridizing conditions, determining the presence of mRNA hybridized to the molecule, and thereby detecting the expression of the mammalian prostate-specific membrane antigen in tissue sections.
- 15 14. An isolated mammalian nucleic acid molecule of claim 2 operatively linked to a promoter of RNA transcription.
- 20 15. A vector which comprises the isolated mammalian nucleic acid molecule of claim 1.
- 25 16. A plasmid of claim 15.
- 30 17. The plasmid of claim 16 designated P55A-PSM (ATCC Accession No. 75294).
- 35 18. A host vector system for the production of a polypeptide having the biological activity of a mammalian prostate-specific membrane antigen which comprises the vector of claim 15 and a suitable host.

-128-

19. A host vector system of claim 18, wherein the suitable host is a bacterial cell, insect cell, or mammalian cell.
- 5 20. A method of producing a polypeptide having the biological activity of a mammalian prostate-specific membrane antigen which comprises growing the host cells of the host vector system of claim 19 under suitable conditions permitting production of the polypeptide and recovering the polypeptide so produced.
- 10 21. A mammalian cell comprising the vector of claim 15.
- 15 22. A method for determining whether a ligand can bind to a mammalian prostate-specific membrane antigen which comprises contacting a mammalian cell having an isolated DNA molecule encoding a mammalian prostate-specific membrane antigen with the ligand under conditions permitting binding of ligands to the mammalian prostate-specific membrane antigen, and determining whether the ligand binds to a mammalian prostate-specific membrane antigen.
- 20 23. A ligand detected by the method of claim 22.
- 25 24. Purified mammalian prostate-specific membrane antigen.
- 30 25. A polypeptide encoded by the isolated mammalian nucleic acid molecule of claim 1.
- 35 26. A method to identify or purify ligands of a mammalian prostate-specific membrane antigen comprising steps of:

-129-

- 5           a)    coupling the purified mammalian prostate-specific membrane antigen of claim 24 to a solid matrix;
- b)    incubating the coupled purified mammalian prostate-specific membrane protein derived from a) with potential ligands under the conditions permitting binding of ligands to the coupled purified mammalian prostate-specific membrane antigen to form a complex;
- 10           c)    washing the ligand and coupled purified mammalian prostate-specific membrane antigen complex formed in b) to eliminate impurities; and
- d)    eluting the ligand from the coupled purified mammalian prostatic membrane specific antigen.
- 15
27.    A ligand identified or purified by claim 26.
28.    A therapeutic agent comprising a ligand of claim 23 or 27 and a cytotoxic agent conjugated thereto.
- 20
29.    The therapeutic agent of claim 28, wherein the cytotoxic agent is a radioisotope or toxin.
- 25       30.    A method of imaging prostate cancer in human patients which comprises administering to the patients at least one ligand of claim 27 or claim 23, capable of binding to the cell surface of the prostate cancer cell and labelled with an imaging agent under conditions permitting binding between the ligand and the cell surface prostate-specific membrane antigen.
- 30
31.    A composition comprising an effective imaging amount the ligand of claim 27 or claim 23 and a
- 35

-130-

pharmaceutically acceptable carrier.

- 5
32. A method to produce antibody using the prostate-specific membrane antigen of claims 24 or 25.
33. A method to produce monoclonal antibody using the mammalian prostate-specific membrane antigen of claims 24 or 25.
- 10 34. An antibody directed against the amino acid sequence of a mammalian prostate-specific membrane antigen.
- 15 35. An antibody directed either to peptide Asp-Glu-Leu-Lys-Ala-Glu (SEQ ID No. 39), or Asn-Glu-Asp-Gly-Asn-Glu (SEQ ID No. 40) or Lys-Ser-Pro-Asp-Glu-Gly (SEQ ID No. 41) of the prostate-specific membrane antigen.
- 20 36. A monoclonal antibody of claim 34 or 35.
37. A therapeutic agent comprising an antibody of claim 34 or 35 and a cytotoxic agent conjugated thereto.
- 25 38. A therapeutic agent of claim 34 or 35, wherein the cytotoxic agent is either a radioisotope or toxin.
- 30 39. A method of imaging prostate cancer in human patient which comprises administering to the patient at least one antibody of claim 34 or 35 capable of binding to the cell surface of the prostate cancer cell and labeled with an imaging agent under conditions permitting formation of a complex between the monoclonal antibody and the
- 35 cell surface prostate-specific membrane antigen.

-131-

40. An imaging method of claim 39 wherein multiple antibodies directed against the amino acid sequence of a mammalian prostate-specific membrane antigen, binding to different epitopes and not interfering with the binding of each other, are administered to the patient.
41. A method of claim 39, wherein the imaging agent is a radioisotope.
42. A prostate cancer specific imaging agent comprising the antibody of claim 34 or 35 and a radioisotope conjugated thereto.
43. A composition comprising an effective imaging amount of the antibody of claim 34 or 35 and a pharmaceutically acceptable carrier.
44. An immunoassay for measuring the amount of a mammalian prostate-specific membrane antigen in a biological sample comprising steps of:
- a) contacting the biological sample with at least one antibody of claim 34 or 35 to form a complex with said antibody and the mammalian prostate-specific membrane antigen, and
  - b) measuring the amount of the prostate-specific membrane antigen in said biological sample by measuring the amount of said complex.
45. An immunoassay for measuring the amount of a mammalian prostate-specific membrane antigen in a biological sample comprising steps of:
- a) contacting the biological sample with at least one ligand of claim 23 or 27 to form a complex with

-132-

said ligand and the mammalian prostate-specific membrane antigen, and

b) measuring the amount of the mammalian prostate-specific membrane antigen in said biological sample by measuring the amount of said complex.

46. A method to purify mammalian prostate-specific membrane antigen comprising steps of:

a) coupling at least one antibody of claim 34 or 35 to a solid matrix;

b) incubating the coupled antibody of a) with a cell lysate containing prostate-specific membrane antigen under the condition permitting binding of the coupled antibody and prostate-specific membrane antigen;

c) washing the solid matrix to eliminate impurities and

d) eluting the prostate-specific membrane antigen from the coupled antibody.

47. A transgenic nonhuman mammal which comprises the isolated nucleic acid molecule of claim 1.

48. A transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a mammalian prostate-specific membrane antigen so placed as to be transcribed into antisense mRNA complementary to mRNA encoding the prostate-specific membrane antigen and which hybridizes to mRNA encoding the mammalian prostate-specific membrane antigen thereby reducing its translation.

49. A method of suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells,

-133-

comprising introducing a DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor cell of a subject, in a way that expression of the prostate specific membrane antigen is under the control of the regulatory element, thereby suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells.

50. The method of claim 49, wherein the DNA molecule encoding prostate specific membrane antigen operatively linked to a 5' regulatory element forms part of a transfer vector which is inserted into a cell or organism, and which vector is capable of replication and expression of prostate specific membrane antigen.
51. The method of claim 50, wherein the DNA molecule encoding prostate specific membrane antigen is integrated into a genome of a eukaryotic or prokaryotic cell.
52. The method of claim 51, wherein a host cell contains and/or expresses a prostate specific membrane antigen.
53. The method of claim 49, wherein the DNA molecule encoding prostate specific membrane antigen is introduced by a delivery vehicle.
54. The method of claim 53, wherein the delivery vehicle is bacterial, viral, fungal, animal, or liposomal.

-134-

55. The method of claim 49, wherein the 5' regulatory element is a promoter.

5 56. The method of claim 49, wherein the 5' regulatory element is an enhancer.

57. The method of claim 49, wherein the subject is a mammal.

10 58. The method of claim 57, wherein the subject is a human.

15 59. A method of suppressing or modulating metastatic ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells, comprising introducing a DNA molecule encoding a prostate specific membrane antigen operatively linked to a 5' regulatory element coupled with a therapeutic DNA into a tumor cell of a subject, thereby suppressing or modulating metastatic  
20 ability of prostate tumor cells, prostate tumor growth or elimination of prostate tumor cells.

25 60. The method of claim 59, wherein the DNA molecule encoding prostate specific membrane antigen operatively linked to a 5' regulatory element forms part of a transfer vector which is inserted into a cell or organism, and which vector is capable of replication and expression of prostate specific  
30 membrane antigen.

35 61. The method of claim 60, wherein the DNA molecule encoding prostate specific membrane antigen is integrated into a genome of a eukaryotic or prokaryotic cell.



-135-

62. The method of claim 61, wherein a host cell contains and/or expresses a prostate specific membrane antigen.

5 63. The method of claim 59, wherein the DNA molecule encoding prostate specific membrane antigen is introduced by a delivery vehicle.

10 64. The method of claim 63, wherein the delivery vehicle is bacterial, viral, fungal, animal, or liposomal.

15 65. The method of claim 59, wherein the 5' regulatory element is a promoter.

66. The method of claim 59, wherein the 5' regulatory element is an enhancer.

20 67. The method of claim 59, wherein the therapeutic DNA encodes a cytokine, viral antigen, or a pro-drug activating enzyme.

25 68. The method of claim 67, wherein the cytokine is interleukin-2.

69. The method of claim 67, wherein the cytokine is interleukin-12.

30 70. The method of claim 67, wherein the cytokine is interferon.

35 71. The method of claim 67, wherein the cytokine is granulocytic macrophage - colony stimulating factor.

-136-

72. The method of claim 59, wherein the subject is a mammal.
73. The method of claim 72, wherein the subject is a human.
74. A prostate tumor cell, comprising a DNA molecule of claim 2 under the control of a prostate specific membrane antigen operatively linked to a 5' regulatory element.
75. The method of claim 74, wherein the 5' regulatory element is a promoter.
76. The method of claim 74, wherein the 5' regulatory element is an enhancer.
77. A therapeutic vaccine for preventing human prostate tumor growth or stimulation of prostate tumor cells in a subject, comprising administering an effective amount to the prostate cell of claim 74, and a pharmaceutical acceptable carrier, thereby preventing the tumor growth or stimulation of tumor cells in the subject.
78. A method of detecting hematogenous micrometastatic tumor cells of a subject, comprising (A) performing nested polymerase chain reaction (PCR) on blood, bone marrow or lymph node samples of the subject using the prostate specific membrane antigen primers, and (B) verifying micrometastases by DNA sequencing and Southern analysis, thereby detecting hematogenous micrometastatic tumor cells of the subject.

-137-

79. The method of claim 78, wherein the primers are derived from prostate specific antigen.
80. The method of claim 78, wherein the subjects is administered hormones in an effective amount.
81. The method of claim 80, wherein the hormone is an androgen suppressor.
82. The method of claim 78, wherein the subject is a mammal.
83. The method of claim 82, wherein the subject is a human.
84. A method of abrogating mitogenic response due to transferrin, comprising introducing a DNA molecule encoding prostate specific membrane antigen operatively linked to a 5' regulatory element into a tumor cell, the expression of which gene is directly associated with a defined pathological effect within a multicellular organism, thereby abrogating mitogen response due to transferrin.
85. The method of claim 84, wherein the 5' regulatory element is a promoter.
86. The method of claim 84, wherein the 5' regulatory element is an enhancer.
87. The method of claim 84, wherein the DNA molecule encoding prostate specific membrane antigen operatively linked to a 5' regulatory element forms part of a transfer vector which is inserted into a cell or organism, and which vector is capable or

-138-


replication and expression of prostate specific membrane antigen.

5        88.    The method of claim 84, wherein the DNA molecule encoding prostate specific membrane antigen is integrated into a genome of a eukaryotic or prokaryotic cell.

10       89.    The method of claim 88, wherein a host cell contains and/or expresses a prostate specific membrane antigen.

1/48

FIGURE 1



2/48

FIGURE 2A




FIGURE 2B





FIGURE 2C




FIGURE 2D




3/48


**FIGURE 3A**



**FIGURE 3B**



**FIGURE 3C**




**FIGURE 3D**



4/48


**FIGURE 4**






5/48

FIGURE 5




6/48

**FIGURE 6A**    **FIGURE 6B**




7/48

FIGURE 7




8/48

FIGURE 8




9/48

FIGURE 9




10/48

**FIGURE 10**



11/48

FIGURE 11



12/48

FIGURE 12A






FIGURE 12B





13/48

FIGURE 13





15/48

FIGURE 14-2

91	H	H	H	H	E	H	C	E	E	E	T	T	C	C	H	H	H	H	H	H	E	E	E	E	E	T	
121	T	T	C	C	E	E	E	E	E	E	C	T	C	C	H	E	E	E	E	E	T	C	C	C	C	T	T
151	T	E	E	E	E	E	E	E	E	E	T	E	C	C	T	C	C	C	C	E	E	E	E	E	E	E	H
181	H	H	H	H	H	H	H	H	H	H	H	H	H	T	T	T	E	E	E	E	E	E	E	E	E	E	E
211	T	T	C	C	H	H	H	H	H	H	H	E	E	E	E	E	E	E	T	E	E	E	E	E	E	E	E
241	E	E	T	T	T	E	C	C	T	T	C	E	E	E	E	E	E	E	E	T	T	C	E	E	E	E	E
271	C	C	C	E	E	E	E	H	E	H	E	E	E	E	E	E	E	C	E	E	E	E	E	E	E	E	E
301	H	H	H	H	H	H	E	T	T	C	C	C	T	E	T	T	E	T	E	E	E	E	E	E	E	E	E
331	E	E	E	C	E	C	H	H	H	H	H	E	E	E	C	C	C	C	E	E	E	E	E	E	E	E	E
361	E	E	E	E	E	E	E	E	E	E	E	E	E	E	C	C	C	T	E	E	E	E	T	C	C	C	T
391	C	H	H	E	E	E	H	H	H	H	H	C	C	C	T	T	T	C	C	C	T	E	E	E	E	E	C
421	H	H	H	H	H	H	C	C	C	C	H	H	H	H	H	H	H	H	H	H	H	E	E	E	E	E	E

16/48

FIGURE 14-3

451 E E C S E E T T E E E E E E E E E E H H H H H H H H H H  
481 H C H H H H H H H H H H T T T C C C C T E E E E E E  
511 E E E C C C E E E E E E H H H H H H H H H H T E C T  
541 T E T T T C E E E E E E E E E E H H H H H H H H H H  
571 H H H H E E E E E E E E E E H H H H H H H H H H  
601 H E E  
631 H H H H H H H H H H H H H H H H H H T T C C E E E E  
661 E E E H E E  
691 E E E T C C C T E E E E E E E E E E H H H H H H C C  
721 C H

-----

17/48

FIGURE 14-4

Semi-graphical output.

=====

Symbols used in the semi-graphical representation:

Helical conformation: X	Extended conformation: -
Turn conformation: >	Coil conformation: *

10	20	30	40	50
MWNLLHETDS	AVATARRPRWLCAGALVLAGGFFLLGFLGFWFIKSSNEAT			

XXXXXXXXXXXXX	-----	XXXXXXXXXXXXX
>	>	>
XXXXXXXXXXXXX	-----	XXXXXXXXXXXXX
>	>	>

60	70	80	90	100
NITPKHNMKAFLDELKAENIKKFLYNFTQIPHLAGTEQNFQIAKQIQSQW				

**FIGURE 14-5**

```
--X-XXXXXXXXXX**>--XXXXXXXXXXXXXXX
--X-XXXXXXXXXX**>--XXXXXXXXXXXXXXX
-X-XXXXXXXXXX**>--XXXXXXXXXXXXXXX
```

KEFGDSELAHYDVLLSYPNKTHPNYISINEDGNEIFNTSLFEP PPPG

^> \* \* \* ^ - - - X \* \* ^ - - - XXXXXXXXXXXX \* \* ^>  
 ^> \* \* \* ^ - - - X \* \* ^ - - - XXXXXXXXXXXX \* \* ^>-

YENVSDI VPPFSAFSPQGMPEGLVVVNYARTEDFFKLERDMKINCSGKI



**FIGURE 14-7**

[illegible]

410 | 420 430 440 450  
SFGTLKKEGWRPRRTILFASWDAAEEFGLGSTWEAEEENSRLLOERGVA YI

--XXXXXX\*--  
 --XXXXXX\*--

--XXXXXX\*--  
 --XXXXXX\*--

NADSSIEGNYTLRVDC TPLMYSLVHNLTKELKSPDEGFEGKSLYESWTKK

\*><XXXXXXXXXXXXXX\*><XXXXXXXXXXXXXX\*><XXXXXXXXXXXXXX\*

510 520 530 540 550  
SPSPEFGMPRI SKLGSGNDFEVFFQRLGIASGRARYTKNWTNKFSGY

[illegible]

560 |  
570 |  
580 |  
590 |  
600 |




21/48

FIGURE 14-8

LYHSVYETVELVEKFDPMFKYHLTVAQVRGGMVFELANSIVLPFDCRDY  
 -----XXXXXXXXXX-X-----XXXXX----->XXX  
 -----XXXXXXXXXX-X-----XXXXX----->XXX  
 610           620           630           640           650  
 |           |           |           |           |  
 AVVLRKYADKIYISMKHPQEMKTYSVSFDLSFSAVKNFTEIASKFSERL  
 XXXXXXXXXXXX-----X\*XXXXXXXXX-----XXXXXXXXXXXXXXXXXXXX  
 XXXXXXXXXXXX-----X\*XXXXXXXXX-----XXXXXXXXXXXXXXXXXXXX  
 660           670           680           690           700  
 |           |           |           |           |  
 QDFDKSNPIVLRMMNDQLMCLERAFIDPLGLPDRPFYRHHVIYAPSSHNKY  
 XX>>>\*>-----XXXXXXXXXX-->>\*>----->\*>\*>  
 XX>>>\*>-----XXXXXXXXXX-->>\*>----->\*>\*>  
 710           720           730           740           750  
 |           |           |           |           |  
 AGESFPGIYDALFDIESKVDPSKAWGEVKRQIYVAAFTVQAAETLSEVA  
 ----->-----XXXXXXXXXX----->-----XXXXXXXXXXXXXXXXXX  
 ----->-----XXXXXXXXXX----->----->-----XXXXXXXXXXXXXXXXXX

22/48

FIGURE 15A



23/48

FIGURE 15B

\*\*\*\*\*  
 \* PREDICTION OF ANTIGENIC DETERMINANTS \*  
 \*\*\*\*\*

Done on sequence PMSANTIGEN.

Total number of residues is: 750.

Analysis done on the complete sequence.

The method used is that of Hopp and Woods.

The averaging group length is: 6 amino acids.

-> This is the value recommended by the authors <-

-----

The three highest points of hydrophilicity are:

( 1)	Ah= 1.62	: From	63 to	68	: Asp-Glu-Leu-Lys-Ala-Glu
( 2)	Ah= 1.57	: From	132 to	137	: Asn-Glu-Asp-Gly-Asn-Glu
( 3)	Ah= 1.55	: From	482 to	487	: Lys-Ser-Pro-Asp-Glu-Gly

Ah stands for: Average hydrophilicity.

Note that, on a group of control proteins, only the highest point was in 100% of the cases assigned to a known antigenic group. The second and third point gave a proportion of 33% of incorrect predictions.

## FIGURE 16-1

The best scores are:

The best scores are:		initn	initl	opt
CHKTFFR	G.gallus mRNA for transferrin receptor	203	120	321
RATTRFR	Rat transferrin receptor mRNA, 3' end.	164	164	311
HUMTFRR	Human transferrin receptor mRNA, complete cd	145	145	266

CHKTFER G.gallus mRNA for transferrin receptor 203 120 321  
51.9% identity in 717 nt overlap

**pmsgen**

1020	1030	1040	1050	1060	1070
TGTCACGCGTGGAAATATCCTAAATCTGAATGGTGAGAGACCCCTCTCACACCAGGTTA					

**CHKTFE**

990	1000	1010	1020	1030	1040
TACACTTATCCCATTTCGGACATGCCCACTTGGAACTGGAGACCCCTTACACCCAGGCTT					

```

1080      1090      1100      1110      1120      1130
pmsgen  CCCAGCAAATGAATATGCTTATAGCGGTGGAATTGCAGAGCGTGTGGTCTTCCAAGTAT
        ::      :      :      :      :      :      :      :      :
CHKTFE  CCCTTCGTTCAACCACACCCA---GTTTCCACCAGTTGAATCTTCAGGACTACCCACAT
        1050      1060      1070      1080      1090      1100

```

	1140	1150	1160	1170	1180	1190
pmsgen	TCCTGTTCA	TCCAATTG	GATACTAT	GATGCACA	GAGCTCCT	AGAAAATGGGTGGCTC
	:	:	:	:	:	:
CHKTFE	TGCTGTTCA	GACCATCT	CTAGCAGT	GCAGCCAG	CGCTGTTCA	GCAAAATGGATGGAGA
	:	:	:	:	:	:
	1110	1120	1130	1140	1150	1160

25/48

FIGURE 16-2

```

1200      1210      1220      1230      1240      1250
pmsgen AGCACCAAGATAGCAGCTGGAGAGGAGTCTCAAGTGCCCTACAATGTTGGACCTGG
      : : : : : : : : : : : : : : : : : :
CHKTFE CACATGCTCTGA-AG--GTTGGAAAGGTGCGATCCA---TTCCTGTAAGGT--GAC--AA
      1170      1180      1190      1200      1210

1260      1270      1280      1290      1300      1310
pmsgen CTTTACTGGAAACTTTTCTACACAAAGTCAAGATGCACATCCACTCTACCAATGAAGT
      : : : : : : : : : : : : : : : : : :
CHKTFE CAAAGCAGGAGA----GCCAGA-TAATGGTGAAACTAGATGTGAACAATTCCTCATGAAAGA
      1220      1230      1240      1250      1260

1320      1330      1340      1350      1360      1370
pmsgen GACAAGAAATTTACAATGTGATAGGTACTCTCAGAGGAGCAGTGGAACACAGACATATGT
      : : : : : : : : : : : : : : : : : :
CHKTFE CAGGAAGATTCTGAACATCTTCGGTGCTATCCAGGGATTGGAAGAACCTGATCGGTATGT
      1270      1280      1290      1300      1310      1320

1380      1390      1400      1410      1420      1430
pmsgen CATTCCTGGGAGGTCACCGGGACTCATGGTGTTTGGTGGTATTGACCCCTCAGAGTGAGC
      : : : : : : : : : : : : : : : : : :
CHKTFE TGTGATTGGAGCCCAAGAGAGACTCCTGGGGCCCAAGAGTGAGTAAAGCTGGCAGTGGAAAC
      1330      1340      1350      1360      1370      1380

```



27/48

FIGURE 16-4

```

1680      1690      1700      1710      1720      1730
pmsgen TACAGCTTGGTACACAACCTAACAAAGAGCTGAAAGCCCTGATGAAGGCTTTGAAGGC
      :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::
CHKTFE TATATGCTGCTGGGGAGTATTATGAAGGGGGTGAAGAATCCAGCAGCAGTCTCAGAGAGC
1630      1640      1650      1660      1670      1680

1740      1750      1760      1770      1780      1790
pmsgen AAATCTCTTTATGAAAGTTGGACTAAATAAGTCCTTCCCCAGAGTTCAGTGGCATGCCC
      :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::
CHKTFE -----CTCTATAACAGACTTGGCCCCAGACTGGGTAAAGCAGTTGTTCCCTCTTGGCCTGGA
1690      1700      1710      1720      1730

```





### FIGURE 16-6

	1320	1330	1340	1350	1360	1370
pmsgen	--AAGTGACAAGAA	TTACAATGTGATAGTACTCTCAGAGGAGCAGTGGAACCA	CACAG			
	:::	:::	:::	:::	:::	:::
RATTRF	GAAAGAAACAAGAA	TACTTAACATCTTTGGCGTTATTAAAGGCTATGAGGGAACCA	CCG			
730	740	750	760	770	780	

```

1380      1390      1400      1410      1420      1430
pmsgen ATATGTCATTCTGGGAGGTCACCGGACTCATGGGTGTTTGGTATTGACCCCTCAGAG
      :: : : : : : : : : : : : : : : : : : :
RATTRF CTACATTGTAGTAGGAGCCAGAGACGCTTGGGGCCCTGGT-GTTCGGAAGTCCAGTG
790      800      810      820      830      840

```

```

1440      1450      1460      1470      1480
pmsgen  T-GGAGCAGCTGTTTCATGAAATTGTGAGGAGCTTTGGAACA-CTGA---AAAAGGAA
:  ::  ::  :  :  :  ::  :  :  :  :  :  :  :  :  :  :  :  :  :  :  :  :
RATTRF  TGGGAACAGGTCTT-CTGTTGAAACTTGCCCAAGTATTCTCAGATATGATTCAAAGAT
850      860      870      880      890      900

```

```

1490      1500      1510      1520      1530      1540
pmsgen  GGTGGAGACCTAGAAGACAATTTGTTTGCAGCTGGGATGCAGAGAAATTTGGTCTT
        :: : X:::: :: :: : :::: ::::: ::::: ::::: ::::: :::::
RATTRF  GGATTAGACCCAGCAGGAGTATTATCTTTGCCAGCTGGACTGCAGGAGACTATGGAGCT
910      920      930      940      950      960

```

## FIGURE 16-7

	1550	1560	1570	1580	1590	1600
pmsgen	CTTGGTTCTACTGAGTGGGCACAGGAGAA---TTCAAGACTCCTTCAAGAGCGTGCCGTG					
RATTRF	GTGGTC CGACTGAGTGGCTGGAGGGGTACCTTTCATCTTTGCATCTAAAG---					
	970	980	990	1000	1010	1020

30/48

	1610	1620	1630	1640	1650	1660
pmsgen	GCTATATTAA	TGCTGACTCATCTA	TAGAGGAAC	TA-CAC	TCTGAGAGT	TGATTGTAC
	:::: :::::	::: : :	: : ::::	: : :::	: : :::	: : :
RATTRF	ACTTACATTTAA	T-CTGGATAAAGTC	GCCTGGGTACT	AGCAACTTCA	AGGTTTCTGCCAG	
	1030	1040	1050	1060	1070	1080

	1670	1680	1690	1700	1710	1720
pmsgen	ACCGCTGATGTACAGCTTGGTACACAACCTAACAAAGAGCTGAAAGC-CCTGATGAAG					
	:: :: : :: :	:	:: :: :	:	:: :: :	:
RATTRF	CCCCCTATTATATACACTTATGGGGAAGATAATGCAGGA--CGTAAAGCATCCGA-----					
	1090	1100	1110	1120	1130	



## FIGURE 16-9

HUMTFRR	Human transferrin receptor mRNA, complete cd	145	145	266
54.3%	identity in 464 nt overlap			

	1230	1240	1250	1260	1270
pmsgen	AGGAAGTCTCAAAGTGCCTACAATGTTGGACCTGGCTTTAC-TGGA	AACTTTCTACAC			
			:	:	:
HUMTFR	TATGGAAGGAGACTGTCCCTCTGACTGGAAAACAGACTCTACATGTAGGATGGTAACCTC				
1140	1150	1160	1170	1180	1190

```

1280      1290      1300      1310      1320      1330
pmsgen AAAAGTCAAGATGCACATC-CACTCT-ACCAATG-----AAGTGACAAGAATTACAA
      : : : : : : : : : : : : : : : : : : : : : : : :
HUMTFR AGAAAGCAAGAATGTGAAGCTCACTGTGAGCAAATGTGCTGAAAGAGATAAAATCTTAA
1200      1210      1220      1230      1240      1250

```

	1340	1350	1360	1370	1380	1390
pmgsen	TGTGATAGTACTCTCAGAGGAGCAGTGGAAC	CAGACAGATATGTCATTCTGGGAGGTCA				
	:	:	:	:	:	:
HUMTER	CATCTTTGGAGTTATTAAAGGCTTTGTAGA	ACCAGATCACTATGTTGTAGTTGGGCCCA				
	:	:	:	:	:	:
	1260	1270	1280	1290	1300	1310

	1400	1410	1420	1430	1440	1450
pmsgen	CCGGACTCATGGTGT	TGTTGGTGAT	TGACCTCAGAGT	-GGAGCAGCTGT	GTTGTTTCATG	
	:	:	:	:	:	:
HUMTFR	GAGAGATGCATGGG	CCCCCTGGAGCT	GC AAAATC	-CGGTGTAGGC	CAGCTCTCCTATTGA	
	1320	1330	1340	1350	1360	1370

33/48

FIGURE 16-10

```

1460      1470      1480      1490      1500
pmsgen AAATTG---TGAGGAGCTTTGGAACACTGAAAAAGGAAGGTTGGAGACCTAGAAACA
:: :: :: :: :: :: :: :: :: X:: :: :: :: :: :: ::
HUMTFR AACTTGCCAGATGTTCTCAGATATGGTCTTAAAGATGGGTTTCAGCCCAGCAGAAGCA
1380      1390      1400      1410      1420      1430

1510      1520      1530      1540      1550      1560
pmsgen TTTTGTGCAAGCTGGGATGCAGAAATTTGGTCTTCTTGGTTCTACTGAGTGGCAG
:: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::
HUMTFR TTATCTTTGCCAGTTGGAGTGTGGAGACTTTGGATCGGTTGGTGCCACTGAATGGCTAG
1440      1450      1460      1470      1480      1490

1570      1580      1590      1600      1610      1620
pmsgen A-GGAGAAATTCAAGACTCCTTCAAGAGCGTGGCGTGGCTTATATTAATGCTGACTCATCT
: :: :: :: :: :: :: :: :: :: :: :: :: :: :: X :: ::
HUMTFR AGGGATACCTTTCGTC-CCTGCATTTAAAGGCTTTCACTTATATTAATCTGGATAAAGCG
1500      1510      1520      1530      1540      1550

1630      1640      1650      1660      1670      1680
pmsgen ATAGAAGGAACACTACACTCTGAGAGTTGATTTGTACACCGCTGATGTACA-GCTTGGT-AC
: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: :: ::
HUMTFR GTTCTTGGTACCAGCAACTCAAGGTTTCTGCCAGCCCACTGTTGTATACGCTTATTGAG
1560      1570      1580      1590      1600      1610

```

## FIGURE 16-11


```

1690      1700      1710      1720      1730      1740
pmsgen  ACAACCTAACAAAGAGCTGAAAGCCCTGATGAAGGCTTTGAAGGCAAATCTCTTATG
          :  ::  :  ::  :
HUMTFR  AAAACAATGCAAAATGTGAAGCATCCGGTTACTGGGCAATTTCTATATCAGGACAGCAAC
1620      1630      1640      1650      1660      1670


```

35/48


**FIGURE 17A**



**FIGURE 17B**




**FIGURE 17C**



36/48


FIGURE 18






37/48

FIGURE 19




38/48

FIGURE 20




39/48

FIGURE 21



40/48

FIGURE 22



**41/48**  
**FIGURE 23**

CELL LINE/TYPE	11p11.2-13 REGION	METASTATIC	PSM RNA DETECTED	PSM DNA DETECTED
LNCap			++	ND
HUMAN PROSTATE			++	ND
A9 (FIBROSARCOMA)	NO	NO	-	-
A9(11) (A9+HUM. 11)	YES	NO	-	REPEAT
AT6.1 (RAT PROSTATE)	NO	YES	-	-
AT6.1-11-c11	YES	NO	+	++
AT6.1-11-c12	NO	YES	-	-
R1564 (RAT MAMMARY)	NO	YES	-	-
R1564-11-c14	YES	YES	-	+
R1564-11-c15	YES	YES	-	REPEAT
R1564-11-c16	YES	YES	-	ND
R1564-11-c12	YES	YES	ND	+

42/48

FIGURE 24A





FIGURE 24B



43/48

FIGURE 25A






FIGURE 25B



44/48


FIGURE 26






45/48

FIGURE 27




46/48

FIGURE 28



47/48

FIGURE 29



**48/48**  
**FIGURE 30**

Patient	Stage	Treatment	PSA	PAP	PSA-PCR	PSM-PCR
1	T2NxMo	None	8.9	0.7	-	+
2	T2NoMo	RRP 7/93	6.1	-	-	+
3	T2CNoMo	PLND 5/93	4.5	0.1	-	+
4	T2BNoMo	RRP 3/92	NMA	0.4	-	+
5	T3NxMo	Proscar + Flutamide	51.3	1.0	-	+
6	Recur T3	I-125 1986	54.7	1.4	-	+
7	T3ANoMo	RRP 10/92	NMA	0.3	-	+
8	T3NxMo	XRT 1987	7.5	0.1	-	-
9	T3NxMo	Proscar + Flutamide	35.4	0.7	-	-
10	D2	S/P XRT Flutamide + Emcyt	311	4.5	+	+
11	D2	RRP 4/91 Lupron 10/92 Velban + Emcyt 12/92	1534	1.4	+	+
12	T2NoMo	RRP 8/91	NMA	0.5	-	+
13	T3NoMo	RRP 1/88 Lupron + Flutamide 5/92	0.1	0.3	-	-
14	D1	PLND 1989 XRT 1989	1.6	0.4	-	-
15	D1	Proscar + Flutamide	20.8	0.5	-	-
16	T2CNoMo	RRP 4/92	0.1	0.3	-	-

SUBSTITUTE SHEET (RULE 26)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10624

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/85.1, 85.8, 88, 93U; 435/6, 7.1, 69.3, 91.2; 514/44; 530/350, 388.1, 389.7; 800/2, DIG1, DIG2; 935/3, 8, 12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	BIOCHIMICA ET BIOPHYSICA ACTA, Volume 1048, issued 1990, T. Solin et al., "Gene Expression and Prostate Specificity of Human Prostatic Acid Phosphatase (PAP): Evaluation by RNA Blot Analyses", pages 72-77, see entire document.	<u>1,4-6,8-9,11,14</u> 12, 13
X	JOURNAL OF BIOLOGICAL CHEMISTRY, Volume 266, Number 17, issued 15 June 1991, P.W. Faber et al., "Characterization of the Human Androgen Receptor Transcription Unit", pages 10743-10749, see entire document.	1-11, 14

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

17 DECEMBER 1993

Date of mailing of the international search report

27 JAN 1994

 Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT  
 Washington, D.C. 20231

Authorized officer

ANTHONY C. CAPUTA

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10624

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	PROCEEDINGS OF THE 82ND ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, Volume 32, issued March 1991, Q. Feng et al., "Purification and Biochemical Characterization of the 7E11-C5 Prostate Carcinoma-Associated Antigen", page 239, see abstract.	<u>24, 25, 34, 36</u> 1-23, 26- 33, 35, 37-89
X Y	CANCER RESEARCH, Volume 50, issued 01 October 1991, A.D. Lopes et al., "Immunohistochemical and Pharmacokinetic Characterization of the Site-Specific Immunoconjugate CYT-356 Derived from Antiprostata Monoclonal Antibody 7E11-C5", pages 6423-6429, see entire document.	<u>34, 36, 44</u> 1-33, 35, 37-43, 45-89
Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, Volume 80, issued March 1983, R.A. Young et al., "Efficient Isolation of Genes by Using Antibody Probes", pages 1194-1198, see entire document.	1-20, 47-89
Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, Volume 88, issued September 1991, B.E. Huber et al., "Retroviral-Mediated Gene Therapy for the Treatment of Hepatocellular Carcinoma: An Innovative Approach For Cancer Therapy", pages 8039-8043, see entire document.	47-77, 80-89
Y	CANCER RESEARCH Volume 51, issued 15 March 1991, T. Mukhopadhyay et al., "Specific Inhibition of K-ras Expression and Tumorigenicity of Lung Cancer Cells by Antisense RNA, pages 1744-1748, see entire document.	47, 48
Y	EUROPEAN JOURNAL OF CANCER, Volume 27, Number 1, issued 1991, M.F. Fey et al., "The Polymerase Chain Reaction: A New Tool for the Detection of Minimal Residual Disease in Hematological Malignancies", pages 89-94, see entire document.	78, 79
Y	US, A, 4,554,101 (HOPP) 19 November 1985, see entire document.	35-38
Y	SCIENCE, Volume 256, issued 12 June 1992, K.W. Culver et al., "In Vivo Gene Transfer with Retroviral Vector-Producer Cells for Treatment of Experimental Brain Tumors", pages 1550-1552, see entire document.	47-77, 80-89

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10624

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	G.J. Tortora et al., "MICROBIOLOGY, AN INTRODUCTION", published 1989 by BENJAMIN/CUMMINGS PUBLISHING CO., INC. (CALIFORNIA), pages 423-426, and 471, see entire document.	28, 29, 32, 33, 35-38
Y	D.P. STITES et al., "BASIC AND CLINICAL IMMUNOLOGY", published 1991 by APPLETON & LANGE (CONNECTICUT), pages 229-251, see entire document.	22, 23, 26-31, 39-43
Y	N.R. ROSE et al., "MANUAL OF CLINICAL LABORATORY IMMUNOLOGY", published 1986 by AMERICAN SOCIETY FOR MICROBIOLOGY (D.C.), pages 88-109, see entire document.	44, 45
Y	W.E. PAUL, "FUNDAMENTAL IMMUNOLOGY", published 1989 by RAVEN PRESS (N.Y. ), pages 628, 629, and 647-651, see entire document.	70, 71
Y	J. SAMBROOK et al., "MOLECULAR CLONING, A LABORATORY MANUAL", published 1989 by COLD SPRING HARBOR LABORATORY PRESS (N.Y.), pages 16.1-16.81, see entire document.	1-21, 47-89
P,Y	CANCER RESEARCH, Volume 53, issued 01 March 1993, R.G. Vile et al., " <u>In Vitro</u> and <u>in Vivo</u> Targeting of Gene Expression to Melanoma Cells", pages 962-967, see entire document.	47-77, 80-89
Y	JOURNAL OF UROLOGY, Volume 143, issued February 1990, H.N. Keer et al., "Elevated Transferrin Receptor Content in Human Prostate Cancer Cell Lines Assessed <u>in Vitro</u> and <u>in Vivo</u> ", pages 381-385, see entire document.	84-89
Y	CELLULAR IMMUNOLOGY, Volume 143, Number 1, issued August 1992, M.K. Gately et al., "Regulation of Human Cytolytic Lymphocyte Responses by Interleukin-12", pages 127-142, see entire document.	68, 69
Y	EUROPEAN JOURNAL OF CANCER, Volume 27, Number 9, issued 1991, A. Decensi et al., "Phase II Study of the Pure Non-steroidal Antiandrogen Nilutamide in Prostatic Cancer", pages 1100-1104, see entire document.	81

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10624

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

A61K 39/00, 39/395, 48/00; C07K 3/12, 15/06, 15/28; C12N 15/12; C12P 19/34; C12Q 1/68; G01N 33/53, 37/00

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/85.1, 85.8, 88, 93U; 435/6, 7.1, 69.3, 91.2; 514/44; 530/350, 388.1, 389.7; 800/2, DIG1, DIG2; 935/3, 8, 12

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

CA, BIOSIS, EMBASE, MEDLINE, PASCAL, DERWENT WORLD PATENTS INDEX, DERWENT BIOTECHNOLOGY ABS, CURRENT BIOTECHNOLOGY ABS, APS  
SEARCH TERMS: PROSTATE, MEMBRANE, ANTIGEN, PSM, 100, ISRAELI, HESTON, FAIR, androgen, IL-12, interleukin 12